

5

THERAPEUTIC APPLICATIONS OF LAMININ AND LAMININ-DERIVED PROTEIN FRAGMENTS

This is a continuation of US Application No. 08/947,057 filed 10/08/1997, which claims priority to US Provisional Application No. 60/027,981 filed 10/08/1996.

10

TECHNICAL FIELD

The invention relates to the discovery, identification and use of laminin, laminin-derived protein fragments, and laminin-derived polypeptides, as well as related peptides and antibodies, for the therapeutic intervention and diagnosis of Alzheimer's disease and other amyloidoses. In addition, the discovery and identification of an Alzheimer's beta-amyloid protein (A β) specific binding region within the globular domain repeats of the laminin A chain, has led to new diagnostic and therapeutic applications for Alzheimer's disease and other amyloidoses which are disclosed.

BACKGROUND OF THE INVENTION

Alzheimer's disease is characterized by the accumulation of a 39-43 amino acid peptide termed the beta-amyloid protein or A β , in a fibrillar form, existing as extracellular amyloid plaques and as amyloid within the walls of cerebral blood vessels. Fibrillar A β amyloid deposition in Alzheimer's disease is believed to be detrimental to the patient and eventually leads to toxicity and neuronal cell death, characteristic hallmarks of Alzheimer's disease. Accumulating evidence now implicates amyloid as a major causative factor of Alzheimer's disease pathogenesis. Discovery and identification of new compounds, agents, proteins, polypeptides or protein-derivatives as potential therapeutic agents to arrest Alzheimer's disease A β amyloid formation, deposition, accumulation and/or persistence is desperately sought.

It is known that A β is normally present in human blood and cerebrospinal fluid.

However, it is not known why this potential fibrillar protein remains soluble in circulating biological fluids. Can the agent(s) responsible for this extraordinary solubility of fibrillar A β be applied to diagnostic and therapeutic regimens against the fibrillar A β amyloid present in Alzheimer's brain?

5

SUMMARY OF THE INVENTION

The present invention provides answers to these questions and relates to the novel and surprising discovery that laminin and specific laminin-derived protein fragments are indeed potent inhibitors of Alzheimer's disease amyloidosis, and therefore have potential use for the therapeutic intervention and diagnosis of the amyloidoses. In addition, we have identified a specific region within laminin which interacts with the Alzheimer's disease beta-amyloid protein and contributes to the observed inhibitory and therapeutic effects. In addition, specific laminin-derived protein fragments which also interact with the A β of Alzheimer's disease have been discovered to be present in human serum and cerebrospinal fluid, and implicate diagnostic applications which are described.

Laminin is a specific basement membrane component that is involved in several fundamental biological processes, and may play important roles in the pathogenesis of a number of different human diseases. Using a solid phase binding immunoassay, the present invention determined that laminin binds the A β of Alzheimer's disease with a single binding constant of $K_d = 2.7 \times 10^{-9}$ M. In addition, using a Thioflavin T fluorometry assay (which quantitatively determines the amount of fibrillar amyloid formed), the present invention has determined that laminin is surprisingly an extremely potent inhibitor of A β fibril formation. In this latter study, 25 μ M of A β (residues 1-40) was incubated at 37°C for 1 week in the

presence or absence of 100 nM laminin. Laminin was found to significantly ($p<0.001$) inhibit A β (1-40) amyloid fibril formation by 2.9-fold at 1 hour, 4.6-fold at 1 day, 30.6-fold at 3 days and 27.1-fold at 1 week. Other basement membrane components including perlecan, fibronectin and type IV collagen were not effective inhibitors of A β (1-40) fibrillogenesis in comparison to laminin, demonstrating the specificity of the inhibitory effect exhibited by laminin. The inhibitory effects of laminin on A β fibrillogenesis was also found to occur in a dose-dependent manner. In addition, laminin was found to cause dissolution of pre-formed Alzheimer's disease amyloid fibrils in a dose-dependent manner following 4 days of incubation. Laminin was digested with V8, trypsin or elastase to determine small protease-resistant fragments of laminin which still interacted with A β . A ~55 kilodalton (kDa) laminin fragment derived from V8 or elastase digested laminin was found to interact with biotinylated A β (1-40). Amino acid sequencing of the ~55 kDa fragment identified an A β -binding domain within laminin situated within the globular repeats of the laminin A chain.

Intact laminin was found to be present in human serum but not human cerebrospinal fluid, whereas laminin protein fragments ranging from ~120 kDa to ~200 kDa were found to be present in both human serum and cerebrospinal fluid. Of all the laminin protein fragments present in human biological fluids described above, a prominent ~130 kilodalton band was found in human serum and cerebrospinal fluid which primarily interacted with A β as determined by ligand blotting methodology. This ~130 kilodalton laminin fragment is known as the E8 fragment (i.e. generated following elastase digestion of laminin)(Yurchenco and Cheng, J. Biol. Chem. 268:17286-17299, 1993) and is also believed to consist of the globular domains of the laminin A chain. The interaction of specific laminin fragments such as the newly discovered ~130 kDa protein is believed to bind A β in biological fluids and keep it in a soluble state. The present invention describes

the use of laminin, laminin-derived protein fragments, and laminin-derived polypeptides for the therapeutic intervention and diagnosis of Alzheimer's disease and other amyloidoses. In addition, the discovery and identification of a specific Alzheimer's A β -binding region within the globular domain repeats of the laminin A chain, and the discovery of the presence of laminin fragments containing this region in human serum and cerebrospinal fluid, has led to new diagnostic and therapeutic applications for Alzheimer's disease and other amyloidoses.

FEATURES OF THE INVENTION

A primary object of the present invention is to establish new therapeutic methods and diagnostic applications for the amyloid diseases. The amyloid diseases include, but are not limited to, the amyloid associated with Alzheimer's disease and Down's syndrome (wherein the specific amyloid is referred to as beta-amyloid protein or A β), the amyloid associated with chronic inflammation, various forms of malignancy and Familial Mediterranean Fever (wherein the specific amyloid is referred to as AA amyloid or inflammation-associated amyloidosis), the amyloid associated with multiple myeloma and other B-cell dyscrasias (wherein the specific amyloid is referred to as AL amyloid), the amyloid associated with type II diabetes (wherein the specific amyloid is referred to as amylin or islet amyloid), the amyloid associated with the prion diseases including Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, kuru and animal scrapie (wherein the specific amyloid is referred to as PrP amyloid), the amyloid associated with long-term hemodialysis and carpal tunnel syndrome (wherein the specific amyloid is referred to as beta₂-microglobulin amyloid), the amyloid associated with senile cardiac amyloid and Familial Amyloidotic Polyneuropathy (wherein the specific amyloid is referred to as transthyretin or prealbumin), and the amyloid associated with endocrine tumors such as medullary carcinoma of the thyroid (wherein the specific amyloid is referred to as variants of procalcitonin).

A primary object of the present invention is to use laminin, laminin-derived protein fragments and/or laminin-derived polypeptides as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses.

5 "Laminin fragments, laminin-derived fragments, laminin-derived protein fragments and/or laminin-derived polypeptides", may include, but are not limited to, laminin A (or A1) chain, laminin B1 chain, laminin B2 chain, laminin A2 chain (merosin), laminin G1 chain, the globular domain repeats within the laminin A1 chain, SEQ ID NO: 1 (11 amino acid sequence within the mouse laminin A chain), SEQ ID NO: 2 (fourth globular repeat with the mouse laminin A chain), SEQ ID NO: 3 (fourth globular repeat within the human laminin A chain), SEQ ID NO: 4 (mouse laminin A chain), SEQ ID NO: 5 (human laminin A chain), SEQ ID NO: 6 (human laminin B1 chain), SEQ ID NO: 7 (mouse laminin B1 chain), SEQ ID NO: 8 (rat laminin B2 chain), SEQ ID NO: 9 (human laminin B2 chain), SEQ ID NO: 10 (mouse laminin G1 chain), SEQ ID NO: 11 (human laminin G1 chain), and all fragments or combinations thereof.

Yet another object of the present invention is to use conformational dependent proteins, polypeptides, or fragments thereof for the treatment of Alzheimer's disease and other amyloidoses. Such conformational dependent proteins include, but are not limited to, laminin, laminin-derived fragments including laminin A1 chain (SEQ ID NO 4; SEQ ID NO: 5), the globular repeat domains within the laminin A1 chain (SEQ ID NO: 2, SEQ ID NO:3), an 11- amino acid peptide sequence within the globular domain of the laminin A chain (SEQ ID NO:1), laminin B1 chain (SEQ ID NO:6, SEQ ID NO: 7), laminin B2 chain (SEQ ID NO: 8, SEQ ID NO:9), laminin G1 chain (SEQ ID NO: 10, SEQ ID NO: 11) and/or portions thereof.

Yet another aspect of the present invention is to use peptidomimetic compounds modelled from laminin, laminin-derived protein fragments and/or laminin-derived polypeptides, including but not limited to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses.

Yet another object of the present invention is to mimic the 3-dimensional A β -binding site(s) on laminin, laminin-derived protein fragments and/or laminin-derived polypeptides and use these mimics as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses.

Yet a further aspect of the present invention is to use anti-idiotypic antibodies to laminin, laminin-derived protein fragments and/or laminin-derived polypeptides as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses.

Another aspect of the invention is to provide new and novel polyclonal and/or monoclonal peptide antibodies which can be utilized in a number of in vitro assays to specifically detect A β -binding laminin derived protein fragments and/or A β -binding laminin derived polypeptides in human tissues and/or biological fluids. Polyclonal or monoclonal antibodies that are made specifically against a peptide portion or fragment of laminin which interacts with A β can be utilized to detect and quantify amyloid disease specific laminin fragments in human tissues and/or biological fluids. These antibodies can be made by administering the peptides in antigenic form to a suitable host. Polyclonal or monoclonal

antibodies may be prepared by standard techniques known to those skilled in the art.

Another object of the present invention is to use laminin, the A β -binding laminin fragments and/or laminin-derived polypeptides referred to above, for the detection and specific localization of laminin peptides important in the amyloid diseases in human tissues, cells, and/or cell culture using standard immunohistochemical techniques.

Yet another aspect of the present invention is to use antibodies recognizing laminin, any of the A β -binding laminin fragments, and/or laminin-derived polypeptides including, but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, for in vivo labelling; for example, with a radionucleotide, for radioimaging to be utilized for in vivo diagnosis, and/or for in vitro diagnosis.

Yet another aspect of the present invention is to make use of laminin, A β -binding laminin protein fragments and/or A β -binding laminin-derived polypeptides including, but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, as potential therapeutics to inhibit the deposition, formation, and accumulation of fibrillar amyloid in Alzheimer's disease and other amyloidoses (described above), and to enhance the clearance and/or removal of preformed amyloid deposits in brain (for Alzheimer's disease and Down's syndrome amyloidosis) and in systemic organs (for systemic amyloidoses).

Another object of the present invention is to use A β -binding laminin-derived

polypeptides or fragments thereof, in conjunction with polyclonal and/or monoclonal antibodies generated against these peptide fragments, using in vitro assays to detect amyloid disease specific autoantibodies in human biological fluids. Specific assay systems can be utilized to not only detect the presence of autoantibodies against A β -binding laminin-derived protein fragments or polypeptides thereof in biological fluids, but also to monitor the progression of disease by following elevation or diminution of laminin protein fragments and/or laminin-derived polypeptide autoantibody levels.

Another aspect of the invention is to utilize laminin, laminin-derived protein fragments and/or laminin-derived polypeptide antibodies and/or molecular biology probes for the detection of these laminin derivatives in human tissues in the amyloid diseases.

Yet another object of the present invention is to use the laminin-derived protein fragments of the present invention in each of the various therapeutic and diagnostic applications described above. The laminin-derived protein fragments include, but are not limited to, the laminin A1 chain, the globular repeats within the laminin A1 chain, the laminin B1 chain, the laminin B2 chain, the laminin G1 chain, the laminin A2 chain (also known as merosin), and all constituents or variations thereof, including but not limited to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, including peptides which have at least 70% homology to the sequences disclosed herein. Specific laminin-derived protein fragments or peptides as described above may be derived from any species including, but are not limited to, human, murine, bovine, porcine, and/or equine species.

Another object of the invention is to provide polyclonal and/or monoclonal peptide

antibodies which can be utilized in a number of in vitro assays to specifically detect laminin protein fragments in human tissues and/or biological fluids. Polyclonal or monoclonal antibodies made specifically against a peptide portion or fragment of any of the laminin fragments described herein can be utilized to detect and quantify laminin-derived protein fragments in human tissues and/or biological fluids. A preferred embodiment is a polyclonal antibody made to the ~130 kilodalton A β -binding laminin fragment present in human serum and cerebrospinal fluid. These antibodies can be made by isolating and administering the laminin-derived fragments and/or polypeptides in antigenic form to a suitable host. Polyclonal or monoclonal antibodies may be prepared by standard techniques by one skilled in the art.

Yet another object of the present invention is to use laminin-derived fragment antibodies as described herein as a specific indicator for the presence and extent of laminin breakdown in brain by monitoring biological fluids including, but not limited to, cerebrospinal fluid, blood, serum, urine, saliva, sputum, and stool.

Yet another object of the present invention is to use laminin-derived fragment antibodies as described herein as a specific indicator for the presence, extent and/or progression of Alzheimer's disease and/or other brain amyloidoses by monitoring biological fluids including, but not limited to, cerebrospinal fluid, blood, serum, urine, saliva, sputum, and stool.

Yet another object of the present invention is to use laminin-derived fragment antibodies as described herein as a specific indicator for the presence and extent of laminin breakdown in systemic organs by monitoring biological fluids including, but not limited to, cerebrospinal fluid, blood, serum, urine, saliva, sputum, and stool.

Yet another object of the present invention is to use laminin-derived fragment antibodies as described herein as a specific indicator for the presence and extent of amyloidosis in type II diabetes by monitoring biological fluids including, but not limited to, cerebrospinal fluid, blood, serum, urine, saliva, sputum, and stool.

Yet another object of the present invention is to use laminin-derived fragment antibodies as described herein as a specific indicator for the presence and extent of amyloidosis in other systemic amyloidoses by monitoring biological fluids including, but not limited to, cerebrospinal fluid, blood, serum, urine, saliva, sputum, and stool.

Yet another object of the present invention is to make use of peptides or fragments of laminin as described herein, including but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, as potential blocking therapeutics for the interaction of laminin and laminin-derived fragments in a number of biological processes and diseases (such as in Alzheimer's disease and other amyloid diseases described herein).

Yet another object of the invention is to utilize specific laminin-derived fragment antibodies, as described herein, for the detection of these laminin fragments in human tissues in the amyloid diseases.

Another object of the present invention is to use laminin, laminin-derived protein fragments, and laminin-derived polypeptides, as described herein, for the treatment of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and

other amyloidoses.

Another object of the present invention is to use pills, tablets, caplets, soft and hard gelatin capsules, lozenges, sachets, cachets, vegicaps, liquid drops, elixers, suspensions, emulsions, solutions, syrups, tea bags, aerosols (as a solid or in a liquid medium), suppositories, sterile injectable solutions, and sterile packaged powders, which contain laminin, laminin-derived protein fragments, and laminin-derived polypeptides, including, but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, to treat patients with Alzheimer's disease and other amyloidoses.

Yet another object of the present invention is to use laminin, laminin-derived protein fragments, and laminin-derived polypeptides, including, but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, as potent agents which inhibit amyloid formation, amyloid deposition, amyloid accumulation, amyloid persistence, and/or cause a dissolution of pre-formed or pre-deposited amyloid fibrils in Alzheimer's disease, and other amyloidoses.

Yet another object of the present invention is to provide the use of laminin, laminin-derived protein fragments, and laminin-derived polypeptides, as described herein, for inhibition of amyloid formation, deposition, accumulation, and/or persistence, regardless of its clinical setting.

Yet another object of the present invention is to provide compositions and methods

involving administering to a subject a therapeutic dose of laminin, laminin-derived protein fragments, and laminin-derived polypeptides, which inhibit amyloid deposition, including but not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof. Accordingly, the compositions and methods of the invention are useful for inhibiting amyloidosis in disorders in which amyloid deposition occurs. The proteins or polypeptides of the invention can be used therapeutically to treat amyloidosis or can be used prophylactically in a subject susceptible to amyloidosis. The methods of the invention are based, at least in part, in directly inhibiting amyloid fibril formation, and/or causing dissolution of preformed amyloid fibrils.

Yet another object of the present invention is to provide pharmaceutical compositions for treating amyloidosis. The pharmaceutical compositions include a therapeutic compound of the invention in an amount effective to inhibit amyloid deposition and a pharmaceutically acceptable vehicle.

These and other features and advantages of the present invention will become more fully apparent when the following detailed description of the invention is read in conjunction with the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention.

FIGURE 1 is a binding curve demonstrating the binding interaction of EHS laminin

to substrate bound A β (1-40). A single binding site with a $K_d = 2.7 \times 10^{-9}$ M is determined.

FIGURE 2 demonstrates the potent inhibition of A β amyloid fibril formation by laminin as determined by a Thioflavin T fluorometry assay over a 1 week experimental period.

FIGURE 3 compares the potent inhibition of A β amyloid fibril formation by laminin to other basement membrane components including fibronectin, type IV collagen and perlecan. Only laminin is found to have a potent inhibitory effect on A β fibrillogenesis as early as 1 hour after incubation.

FIGURE 4 is a graph of a 1 week Thioflavin T fluorometry assay utilized to determine the potential dose-dependent effects of laminin on inhibition of A β amyloid fibril formation. Significant dose-dependent inhibition of A β (1-40) amyloid fibril formation is observed at 1 day, 3 days and 1 week of treatment with increasing concentrations of laminin.

FIGURE 5 is a graph of a Thioflavin T fluorometry assay utilized to determine the potential dose-dependent effects of laminin on dissolution of pre-formed A β (1-40) amyloid fibrils within a 4 day incubation period. Laminin causes dissolution of pre-formed A β amyloid fibrils in a dose-dependent manner.

FIGURE 6 is a graph of a 1 week Thioflavin T fluorometry assay utilized to determine the effects of laminin on islet amyloid polypeptide (amylin) fibrillogenesis, and determine whether laminin causes a dose-dependent inhibition of amylin fibril formation. Laminin does not significantly inhibit amylin fibrillogenesis suggesting its specificity for

Alzheimer's disease amyloidosis.

FIGURE 7 is a black and white photograph of laminin digested with V8 protease, separated by SDS-PAGE and following interaction with biotinylated A β (1-40). The smallest fragment of V8-resistant laminin that interacts with A β is a ~55 kilodalton fragment.

FIGURE 8 is a black and white photograph of laminin digested with trypsin, separated by SDS-PAGE and following interaction with biotinylated A β (1-40). The smallest fragment of trypsin-resistant laminin that interacts with A β is a ~30 kilodalton fragment.

FIGURE 9 is a black and white photograph of laminin digested with elastase, separated by SDS-PAGE and following interaction with biotinylated A β (1-40). A ~55 kilodalton laminin fragment (arrow) that binds biotinylated A β was identified and sequenced. Note also the presence of a ~130 kDa fragment (arrowheads) that binds A β following 1.5 hours of elastase digestion (lane 2). Panel A is a ligand blot using biotinylated A β as a probe, whereas panel B is Coomassie blue staining of the same blot in Panel A to locate the specific band(s) for sequencing.

FIGURE 10 shows the complete amino acid sequence of the mouse laminin A chain. Sequencing of the ~55 kilodalton A β -binding band shown in Figure 9 leads to the identification of an 11 amino acid segment (underline and arrowhead) within the laminin A chain. This A β binding region of laminin is situated within the globular domain repeats of the laminin A chain.

FIGURE 11 shows schematic diagrams of laminin and the newly discovered "A β -

binding region” of laminin (shown in left panel; between the two arrowheads) which is situated within the last three globular domains of the laminin A chain.

FIGURE 12 is a black and white photograph of a Western blot demonstrating the presence of laminin (arrowheads) and/or laminin-derived protein fragments (bands between the two arrows) in human serum (lanes 1-7; left side) and human cerebrospinal fluid (lanes 1-7; right side) obtained from Alzheimer’s disease, type II diabetes and normal aged patients. A ~110-130 kilodalton range of laminin positive protein fragments (between the two arrows) is present in both human serum and cerebrospinal fluid, whereas intact laminin (arrowheads) is only present in serum but not in cerebrospinal fluid.

FIGURE 13 is a black and white photograph demonstrating that intact laminin (arrow) and a prominent ~130 kilodalton band (arrowhead) present in human Alzheimer’s disease, type II diabetes and normal aged patient serum, bind A β . The A β -binding laminin and specific A β -binding laminin fragments in human serum were identified following separation by SDS-PAGE and interaction with nanomolar concentrations of biotinylated A β (1-40).

FIGURE 14 is a black and white photograph demonstrating the presence of a prominent ~130 kilodalton band (arrow) in human Alzheimer’s disease and normal aged patient cerebrospinal fluid, identified following separation by SDS-PAGE and following interaction with nanomolar concentrations of biotinylated A β (1-40). This same ~130 kilodalton A β -binding protein is also present in human serum (Figure 13).

DETAILED DESCRIPTION OF THE INVENTION

The following sections are provided by way of background to better appreciate the invention.

Alzheimer's Disease

Alzheimer's disease is the most common cause of dementia in middle and late life, and is manifested by progressive impairment of memory, language, visuospatial perceptions and behavior (A Guide to the Understanding of Alzheimer's Disease and Related Disorders, edited by Jorm, New York University Press, New York 1987). A diagnosis of probable Alzheimer's disease can be made on clinical criteria (usually by the exclusion of other diseases, memory tests etc), but a definite diagnosis requires the histological examination of specific abnormalities in the brain tissue usually obtained at autopsy.

In Alzheimer's disease, the parts of the brain essential for cognitive processes such as memory, attention, language, and reasoning degenerate, robbing victims of much that makes us human, including independence. In some inherited forms of Alzheimer's disease, onset is in middle age, but more commonly, symptoms appear from the mid-60's onward. Alzheimer's disease is characterized by the deposition and accumulation of a 39-43 amino acid peptide termed the beta-amyloid protein, A β or β /A4 (Glenner and Wong, Biochem. Biophys. Res. Comm. 120:885-890, 1984; Masters et al, Proc. Natl. Acad. Sci. USA 82:4245-4249, 1985; Husby et al, Bull. WHO 71:105-108, 1993). A β is derived from larger precursor proteins termed beta-amyloid precursor proteins (or β PPs) of which there are several alternatively spliced variants. The most abundant forms of the β PPs include

proteins consisting of 695, 751 and 770 amino acids (Tanzi et al, Nature 331:528-530, 1988; Kitaguchi et al, Nature 331:530-532, 1988; Ponte, et al, Nature 331:525-528, 1988).

The small A β peptide is a major component which makes up the amyloid deposits of neuritic “plaques” and in the walls of blood vessels (known as cerebrovascular amyloid deposits) in the brains of patients with Alzheimer’s disease. In addition, Alzheimer’s disease is characterized by the presence of numerous neurofibrillary “tangles”, consisting of paired helical filaments which abnormally accumulate in the neuronal cytoplasm (Grundke-Iqbal et al, Proc. Natl. Acad. Sci. USA 83:4913-4917, 1986; Kosik et al, Proc. Natl. Acad. Sci. USA 83:4044-4048, 1986; Lee et al, Science 251:675-678, 1991). The pathological hallmarks of Alzheimer’s disease is therefore the presence of “plaques” and “tangles”, with amyloid being deposited in the central core of plaques and within the blood vessel walls. It is important to note that a so-called “normal aged brain” has some amyloid plaques and neurofibrillary tangles present. However, in comparison, an Alzheimer’s disease brain shows an over abundance of plaques and tangles. Therefore, differentiation of an Alzheimer’s disease brain from a normal brain from a diagnostic point of view is primarily based on quantitative assessment of “plaques” and “tangles”.

In an Alzheimer’s disease brain, there are usually thousands of neuritic plaques. The neuritic plaques are made up of extracellular deposits consisting of an amyloid core usually surrounded by enlarged axons and synaptic terminals, known as neurites, and abnormal dendritic processes, as well as variable numbers of infiltrating microglia and surrounding astrocytes. The neurofibrillary tangles present in the Alzheimer’s disease brain mainly consist of tau protein, which is a microtubule-associated protein (Grundke-Iqbal et al, Proc. Natl. Acad. Sci. USA 83:4913-4917, 1986; Kosik et al, Proc. Natl. Acad. Sci. USA 83:4044-4048, 1986; Lee et al, Science 251:675-678, 1991). At the ultrastructural level, the tangle consists of paired helical filaments twisting like a ribbon, with a specific crossing

over periodicity of 80 nanometers. In many instances within a neurofibrillary tangle, there are both paired helical filaments and straight filaments. In addition, the nerve cells will many times die, leaving the filaments behind. These tangles are known as “ghost tangles” since they are the filamentous remnants of the dead neuron.

5

The other major type of lesion found in the brain of an Alzheimer’s disease patient is the accumulation of amyloid in the walls of blood vessels, both within the brain parenchyma and in the walls of the larger meningeal vessels which lie outside the brain. The amyloid deposits localized to the walls of blood vessels are referred to as cerebrovascular amyloid or congophilic angiopathy (Mandybur, J. Neuropath. Exp. Neurol. 45:79-90, 1986; Pardridge et al, J. Neurochem. 49:1394-1401, 1987).

10

In addition, Alzheimer’s disease patients demonstrate neuronal loss and synaptic loss. Furthermore, these patients also exhibit loss of neurotransmitters such as acetylcholine. Tacrine, the first FDA approved drug for Alzheimer’s disease is a cholinesterase inhibitor (Cutler and Sramek, New Engl. J. Med. 328:808-810, 1993). However, this drug has showed limited success, if any, in the cognitive improvement in Alzheimer’s disease patients and initially had major side effects such as liver toxicity.

15

20

For many years there has been an ongoing scientific debate as to the importance of “amyloid” in Alzheimer’s disease and whether the “plaques” and “tangles” characteristic of this disease, were a cause or merely the consequences of the disease. Recent studies during the last few years have now implicated that amyloid is indeed a causative factor for Alzheimer’s disease and not merely an innocent bystander. The Alzheimer’s disease A β protein in cell culture has been shown to cause degeneration of nerve cells within short periods of time (Pike et al, Br. Res. 563:311-314, 1991; J. Neurochem. 64:253-265,

25

1994). Studies suggest that it is the fibrillar structure, a characteristic of all amyloids, that is responsible for the neurotoxic effects. The A β has also been found to be neurotoxic in slice cultures of hippocampus (the major memory region affected in Alzheimer's)(Harrigan et al, Neurobiol. Aging 16:779-789, 1995) and induces nerve cell death in transgenic mice

5 (Games et al, Nature 373:523-527, 1995; Hsiao et al, Neuron 15:1203-1218, 1995). In addition, injection of the Alzheimer's A β into rat brain causes memory impairment and neuronal dysfunction (Flood et al, Proc. Natl. Acad. Sci. U.S.A. 88:3363-3366, 1991; Br. Res. 663:271-276, 1994), two additional hallmarks of Alzheimer's disease. Probably, the most convincing evidence that amyloid (ie. beta-amyloid protein) is directly involved in the
10 pathogenesis of Alzheimer's disease comes from genetic studies. It has been discovered that the production of A β can result from mutations in the gene encoding, its precursor, known as the beta-amyloid precursor protein (Van Broeckhoven et al, Science 248:1120-1122, 1990; Europ. Neurol. 35:8-19, 1995; Murrell et al, Science 254:97-99, 1991; Haass et al, Nature Med. 1:1291-1296, 1995). This precursor protein when normally processed usually
15 only produces very little of the toxic A β . The identification of mutations in the amyloid precursor protein gene which causes familial, early onset Alzheimer's disease is the strongest argument that amyloid is central to the pathogenetic process underlying this disease. Four reported disease-causing mutations have now been discovered which demonstrate the importance of the beta-amyloid protein in causing familial Alzheimer's
20 disease (reviewed in Hardy, Nature Genet. 1:233-234, 1992). All of these studies suggest that providing a drug to reduce, eliminate or prevent fibrillar A β formation, deposition, accumulation and/or persistence in the brains of human patients should be considered an effective therapeutic.

Other Amyloid Diseases

The “amyloid diseases” consist of a group of clinically and generally unrelated human diseases which all demonstrate a marked accumulation in tissues of an insoluble extracellular substance known as “amyloid”, and usually in an amount sufficient to impair normal organ function. Rokitansky in 1842 (Rokitansky, “Handbuch der pathologischen Anatomie”, Vol. 3, Braumuller and Seidel, Vienna) was the first to observe waxy and amorphous looking tissue deposits in a number of tissues from different patients. However, it wasn’t until 1854 when Virchow (Virchow, Arch. Path. Anat. 8:416, 1854) termed these deposits as “amyloid” meaning “starch-like” since they gave a positive staining with the sulfuric acid-iodine reaction, which was used in the 1850’s for demonstrating cellulose. Although cellulose is not a constituent of amyloid, nonetheless, the staining that Virchow observed was probably due to the present of proteoglycans (PGs) which appear to be associated with all types of amyloid deposits. The name amyloid has remained despite the fact that Friederich and Kekule in 1859 discovered the protein nature of amyloid (Friedrich and Kekule, Arch. Path. Anat. Physiol. 16:50, 1859). For many years, based on the fact that all amyloids have the same staining and structural properties, lead to the postulate that a single pathogenetic mechanism was involved in amyloid deposition , and that amyloid deposits were thought to be composed of a single set of constituents. Current research has clearly shown that amyloid is not a uniform deposit and that amyloids may consist of different proteins which are totally unrelated (Glenner, N. England J. Med. 302:1283-1292, 1980).

Although the nature of the amyloid itself has been found to consist of completely different and unrelated proteins, all amyloids appear similar when viewed under the microscope due to amyloid’s underlying protein able to adapt into a fibrillar structure. All

amyloids regardless of the nature of the underlying protein 1) stain characteristically with the Congo red dye and display a classic red/green birefringence when viewed under polarized light (Puchtler et al, J. Histochem. Cytochem. 10:355-364, 1962), 2) ultrastructurally consists of fibrils with a diameter of 7-10 nanometers and of indefinite length, 3) adopt a predominant beta-pleated sheet secondary structure. Thus, amyloid fibrils viewed under an electron microscope (30,000 times magnification) from the post-mortem brain of an Alzheimer's disease patient would look nearly identical to the appearance of amyloid present in a biopsied kidney from a rheumatoid arthritic patient. Both these amyloids would demonstrate a similar fibril diameter of 7-10 nanometers.

In the mid to late 1970's amyloid was clinically classified into 4 groups, primary amyloid, secondary amyloid, familial amyloid and isolated amyloid. Primary amyloid, is amyloid appearing de novo, without any preceding disorder. In 25-40% of these cases, primary amyloid was the antecedent of plasma cell dysfunction such as the development of multiple myeloma or other B-cell type malignancies. Here the amyloid appears before rather than after the overt malignancy. Secondary amyloid, appeared as a complication of a previously existing disorder. 10-15% of patients with multiple myeloma eventually develop amyloid (Hanada et al, J. Histochem. Cytochem. 19:1-15, 1971). Patients with rheumatoid arthritis, osteoarthritis, ankylosing spondylitis can develop secondary amyloidosis as with patients with tuberculosis, lung abscesses and osteomyelitis (Benson and Cohen, Arth. Rheum. 22:36-42, 1979; Kamei et al, Acta Path. Jpn. 32:123-133, 1982; McAdam et al, Lancet 2:572-575, 1975). Intravenous drug users who self-administer and who then develop chronic skin abscesses can also develop secondary amyloid (Novick, Mt. Sin. J. Med. 46:163-167, 1979). Secondary amyloid is also seen in patients with specific malignancies such as Hodgkin's disease and renal cell carcinoma (Husby et al, Cancer Res. 42:1600-1603, 1982). Although these were all initially classified as secondary amyloid,

once the amyloid proteins were isolated and sequenced many of these turned out to contain different amyloid proteins.

The familial forms of amyloid also showed no uniformity in terms of the peptide responsible for the amyloid fibril deposited. Several geographic populations have now been identified with genetically inherited forms of amyloid. One group is found in Israel and this disorder is called Familial Mediterranean Fever and is characterized by amyloid deposition, along with recurrent inflammation and high fever (Mataxas, Kidney 20:676-685, 1981). Another form of inherited amyloid is Familial Amyloidotic Polyneuropathy, and has been found in Swedish (Skinner and Cohen, Biochem. Biophys. Res. Comm. 99:1326-1332, 1981), Portuguese (Saraiva et al, J. Lab. Clin. Med. 102:590-603, 1983; J. Clin. Invest. 74:104-119, 1984) and Japanese (Tawara et al, J. Lab. Clin. Med. 98:811-822, 1981) nationalities. Amyloid deposition in this disease occurs predominantly in the peripheral and autonomic nerves. Hereditary amyloid angiopathy of Icelandic origin is an autosomal dominant form of amyloid deposition primarily affecting the vessels in the brain, and has been identified in a group of families found in Western Iceland (Jennson et al, Clin. Genet. 36:368-377, 1989). These patients clinically have massive cerebral hemorrhages in early life which usually causes death before the age of 40.

The primary, secondary and familial forms of amyloid described above tend to involve many organs of the body including heart, kidney, liver, spleen, gastrointestinal tract, skin, pancreas, and adrenal glands. These amyloid diseases are also referred to as "systemic amyloids" since so many organs within the body demonstrate amyloid accumulation. For most of these amyloidoses, there is no apparent cure or effective treatment and the consequences of amyloid deposition can be detrimental to the patient. For example, amyloid deposition in kidney may lead to renal failure, whereas amyloid

deposition in heart may lead to heart failure. For these patients, amyloid accumulation in systemic organs leads to eventual death generally within 3 to 5 years.

Isolated forms of amyloid, on the other hand, tend to involve a single organ system.

5 Isolated amyloid deposits have been found in the lung, and heart (Wright et al, Lab. Invest. 30:767-773, 1974; Pitkanen et al, Am. J. Path. 117:391-399, 1984). Up to 90% of type II diabetic patients (non-insulin dependent form of diabetes) have isolated amyloid deposits in the pancreas restricted to the beta cells in the islets of Langerhans (Johnson et al, New Engl. J. Med. 321:513-518, 1989; Lab. Invest. 66:522-535, 1992). Isolated forms of amyloid
10 have also been found in endocrine tumors which secrete polypeptide hormones such as in medullary carcinoma of the thyroid (Butler and Khan, Arch. Path. Lab. Med. 110:647-649, 1986; Berger et al, Virch. Arch. A Path. Anat. Hist. 412:543-551, 1988). A serious complication of long term hemodialysis is amyloid deposited in the medial nerve and clinically associated with carpal tunnel syndrome (Gejyo et al, Biochem. Biophys. Res. Comm. 129:701-706, 1985; Kidney Int. 30:385-390, 1986). By far, the most common type
15 and clinically relevant type of organ-specific amyloid, and amyloid in general, is that found in the brains of patients with Alzheimer's disease (see U.S. Patent No. 4,666,829 and Glenner and Wong, Biochem. Biophys. Res. Comm. 120:885-890, 1984; Masters et al, Proc. Natl. Acad. Sci., USA 82:4245-4249, 1985). In this disorder, amyloid is
20 predominantly restricted to the central nervous system. Similar deposition of amyloid in the brain occurs in Down's syndrome patients once they reach the age of 35 years (Rumble et al, New England J. Med. 320:1446-1452, 1989; Mann et al, Neurobiol. Aging 10:397-399, 1989). Other types of central nervous system amyloid deposition include rare but highly infectious disorders known as the prion diseases which include Creutzfeldt-Jakob disease,
25 Gerstmann-Straussler syndrome, and kuru (Gajdusek et al, Science 197:943-960, 1977; Prusiner et al, Cell 38:127-134, 1984; Prusiner, Scientific American 251:50-59, 1984;

Prusiner et al, Micr. Sc. 2:33-39, 1985; Tateishi et al, Ann. Neurol. 24:35-40, 1988).

It was misleading to group the various amyloidotic disorders strictly on the basis of their clinical features, since when the major proteins involved were isolated and sequenced, they turned out to be different. For example, amyloid seen in rheumatoid arthritis and osteoarthritis, now known as AA amyloid, was the same amyloid protein identified in patients with the familial form of amyloid known as Familial Mediterranean Fever. Not to confuse the issue, it was decided that the best classification of amyloid should be according to the major protein found, once it was isolated, sequenced and identified.

Thus, amyloid today is classified according to the specific amyloid protein deposited. The amyloid diseases include, but are not limited to, the amyloid associated with Alzheimer's disease, Down's syndrome and hereditary cerebral hemorrhage with amyloidosis of the Dutch type (wherein the specific amyloid is now known as the beta-amyloid protein or A β), the amyloid associated with chronic inflammation, various forms of malignancy and Familial Mediterranean Fever (AA amyloid or inflammation-associated amyloidosis), the amyloid associated with multiple myeloma and other B-cell abnormalities (AL amyloid), the amyloid associated with type II diabetes (amylin or islet amyloid), the amyloid associated with the prion diseases including Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, kuru and animal scrapie (PrP amyloid), the amyloid associated with long-term hemodialysis and carpal tunnel syndrome (beta₂-microglobulin amyloid), the amyloid associated with senile cardiac amyloid and Familial Amyloidotic Polyneuropathy (prealbumin or transthyretin amyloid), and the amyloid associated with endocrine tumors such as medullary carcinoma of the thyroid (variants of procalcitonin).

Laminin and Its Structural Domains

Laminin is a large and complex 850 kDa glycoprotein which normally resides on the basement membrane and is produced by a variety of cells including embryonic, epithelial and tumor cells (Foidart et al, Lab. Invest. 42:336-342, 1980; Timpl et al, Methods Enzymol. 82:831-838, 1982). Laminin-1 (is derived from the Engelbreth-Holm-Swarm tumor) and is composed of three distinct polypeptide chains, A, B1 and B2 (also referred to as alpha1, B1 and gamma-1, respectively), joined in a multidomain structure possessing three short arms and one long arm (Burgeson et al, Matrix Biol. 14:209-211, 1994). Each of these arms is subdivided into globular and rodlike domains. Studies involving in vitro self-assembly and the analysis of cell-formed basement membranes have shown that laminin exists as a polymer, forming part of a basement membrane network (Yurchenco et al, J. Biol. Chem. 260:7636-7644, 1985; Yurchenco et al, J. Cell Biol. 117:1119-1133, 1992; Yurchenco and Cheng, J. Biol. Chem. 268: 17286-17299, 1993). Laminin is believed to play important roles in a number of fundamental biological processes including promotion of neural crest migration (Newgreen and Thiery, Cell Tissue Res. 211:269-291, 1980; Rovasio et al, J. Cell Biol. 96:462-473, 1983), promotion of neurite outgrowth (Lander et al, Proc. Natl. Acad. Sci. 82:2183-2187, 1985; Bronner-Fraser and Lallier, Cell Biol. 106:1321-1329, 1988), the formation of basement membranes (Kleinman et al, Biochem. 22:4969-4974, 1983), the adhesion of cells (Engvall et al, J. Cell Biol. 103: 2457-2465, 1986) and is inducible in adult brain astrocytes by injury (Liesi et al, EMBO J. 3:683-686, 1984). Laminin interacts with other components including type IV collagen (Terranova et al, Cell 22:719-726, 1980; Rao et al, Biochem. Biophys. Res. Comm. 128:45-52, 1985; Charonis et al, J. Cell Biol. 100: 1848-1853, 1985; Laurie et al, J. Mol. Biol. 189:205-216, 1986), heparan sulfate proteoglycans (Riopelle and Dow, Brain Res. 525:92-100, 1990; Battaglia et al, Eur. J. Biochem. 208:359-366, 1992) and heparin (Sakashita et al,

FEBS Lett. 116:243-246, 1980; Del Rosso et al, Biochem. J. 199:699-704, 1981; Skubitz et al, J. Biol. Chem. 263:4861-4868, 1988).

Several of the functions of laminin have been found to be associated with the short arms. First, the short arms have been found to participate in laminin polymerization (Yurchenco et al, J. Cell Biol. 117:1119-1133, 1992; Yurchenco and Cheng, J. Biol. Chem. 268: 17286-17299, 1993). A recently proposed three-arm interaction hypothesis of laminin polymerization (Yurchenco and Cheng, J. Biol. Chem. 268: 17286-17299, 1993) further holds that self-assembly is mediated through the end regions of each of the three short arms. A prediction of this model is that each short arm can independently and competitively inhibit laminin polymerization. However, it has not been possible to formally test this prediction using conventional biochemical techniques because of an inability to separate the alpha and gamma chains. Second, several heparin binding sites have been thought to reside in the short arms (Yurchenco et al, J. Biol. Chem. 265:3981-3991, 1990; Skubitz et al, J. Cell Biol. 115:1137-1148, 1991), although the location of these sites have remained obscure. Third, the alpha1β1 integrin has been found to selectively interact with large short arm fragments containing all or most of the short arm domains (Hall et al, J. Cell Biol. 110:2175-2184, 1990; Goodman et al, J. Cell Biol. 113:931-941, 1991).

Most functional activities of laminin appear to be dependent upon the conformational state of the glycoprotein. Specifically, self-assembly and its calcium dependence, nidogen (entactin) binding to laminin, alpha6β1 integrin recognition of the long arm, heparin binding to the proximal G domain (cryptic) and RGD-dependent recognition of the short A chain of laminin (cryptic) have all been found to be conformationally dependent (Yurchenco et al, J. Biol. Chem. 260:7636-7644, 1985; Fox et al, EMBO J. 10:3137-3146, 1991; Sung et al, J. Cell Biol. 123:1255-1268, 1993). Two consequences of improperly folded laminin, loss of

normal functional activity and the activation of previously cryptic activities, suggest that it is important to map and characterize biological activities using correctly folded laminin or conformational homologues to any particular laminin or laminin fragment.

5 Laminin may also be involved in the pathogenesis of a number of important diseases. For example, in diabetes significant decrease in the levels of laminin on the glomerular basement membranes indicates that a molecular imbalance occurs (Shimomura and Spiro, Diabetes 36:374-381, 1987). In experimental AA amyloidosis (ie. inflammation-associated amyloidosis), increased levels of laminin are observed at the sites of AA amyloid
10 deposition (Lyon et al, Lab. Invest. 64:785-790, 1991). However, the role(s) of laminin in systemic amyloidosis is not known. In Alzheimer's disease and Down's syndrome, laminin is believed to be present in the vicinity of A β -containing amyloid plaques (Perlmutter and Chui, Brain Res. Bull. 24:677-686, 1990; Murtomaki et al, J. Neurosc. Res. 32:261-273, 1992; Perlmutter et al, Micro. Res. Tech. 28:204-215, 1994).

15 Previous studies have indicated that the various isoforms of the beta-amyloid precursor proteins of Alzheimer's disease, bind both the basement membrane proteins perlecan (Narindrasorasak et al, J. Biol. Chem. 266:12878-12883, 1991) and laminin (Narindrasorasak et al, Lab. Invest. 67:643-652, 1992). With regards to laminin, it was not
20 previously known whether laminin interacts with A β , whether a particular domain of laminin (if any) participates in A β interactions, and whether laminin had any significant role(s) in A β amyloid fibrillogenesis.

25 The present invention has discovered that laminin binds A β with relatively high affinity and surprisingly laminin is a potent inhibitor of A β amyloid formation, and causes dissolution of pre-formed Alzheimer's disease amyloid fibrils. In addition, a 55-kilodalton

elastase resistant fragment of laminin which also binds A β has been localized to the globular domain repeats within the A chain of laminin. This region is believed to be responsible for many of the inhibitory effects that laminin has on Alzheimer's disease amyloidosis. These findings indicate that laminin, laminin-derived protein fragments and/or laminin-derived polypeptides, particularly those containing the disclosed A β -binding site within the globular domain repeats within the laminin A chain, may serve as novel inhibitors of A β amyloidosis in Alzheimer's disease and other amyloidoses. In addition, the discovery and identification of an Alzheimer's A β -binding region within the globular domain repeats of the laminin A chain, and the discovery of its presence in human serum and cerebrospinal fluid, as a ~130 kDa laminin-derived fragment, leads to novel diagnostic and therapeutic applications for Alzheimer's disease and other amyloidoses.

Examples

The following examples are provided to disclose in detail preferred embodiments of the binding interaction of laminin with A β , and the potent inhibitory effects of laminin and disclosed fragments on A β fibril formation. However, it should not be construed that the invention is limited to these specific examples.

Example 1

Binding of Laminin to the Beta-Amyloid Protein (A β) of Alzheimer's Disease

2 μ g of A β (1-40)(Bachem Inc., Torrance, CA USA; Lot #WM365) in 40 μ l of Tris-buffered saline (TBS)(pH 7.0) was allowed to bind overnight at 4°C to microtiter wells (Nunc plates, Maxisorb). The next day all of the microtiter wells were blocked by

incubating with 300 µl of Tris-buffered saline containing 100 mM Tris-HCl, 50 mM NaCl, 0.05% Tween-20, and 3 mM NaN₃ (pH 7.4)(TTBS) plus 2% bovine serum albumin (BSA). Various dilutions (ie. 1:10, 1:30, 1:90, 1:270, 1:810, 1:2430 and 1:7290) of Engelbreth-Holm-Swarm (EHS) mouse tumor laminin (1 mg/ml)(Sigma Chemical Co., St. Louis, MO, USA) in 250 µl of TBS (pH 7.4) were placed in wells (in triplicate) either containing substrate bound Aβ (1-40) or blank, and allowed to bind overnight at 4°C overnight. The next day, the wells were rinsed 3 times with TTBS, and then probed for 2 hours with 100 µl of rabbit anti-laminin antibody (Sigma Chemical Company, St. Louis, MO) diluted 1:10,000 in TTBS. After 3 rinses with TTBS, the wells were then incubated for 2 hours on a rotary shaker with 100 µl of secondary probe consisting of biotinylated goat anti-rabbit (1:1000) and streptavidin-peroxidase (1:500 dilution of a 2 µg/ml solution) in TTBS containing 0.1% BSA. The wells were then rinsed 3 times with TTBS and 100 µl of a substrate solution (OPD-Sigma Fast from Sigma Chemical Co., St. Louis, MO) was added to each well and allowed to develop for 10 minutes or until significant color differences were observed. The reaction was stopped with 50 µl of 4N H₂SO₄ and read on a Model 450 microplate reader (Biorad, Hercules, CA USA) at 490 nm. Data points representing a mean of triplicate determinations were plotted and the affinity constants (ie. K_d) were determined using Ultrafit (version 2.1, Biosoft, Cambridge, U.K.) as described below.

The binding data were analyzed assuming a thermodynamic equilibrium for the formation of the complex BL, from the laminin ligand in solution, L, and the uncomplexed Aβ adsorbed to the microtiter well, B, according to the equation: $K_d = [B] \times [L] / [BL]$. We elected to determine K_d's by using an enzyme-linked immunoassay that gives a color signal that is proportional to the amount of unmodified laminin bound to Aβ (Engel, J. and

Schalch, W., Mol. Immunol. 17:675-680, 1980; Mann, K. et al, Eur. J. Biochem. 178:71-80, 1988; Fox, J.W. et al, EMBO J. 10:3137-3146, 1991; Battaglia, C. et al, Eur. J. Biochem. 208:359-366, 1992).

5 To account for potential non-specific binding, control wells without A β (in triplicate) were included for each concentration of laminin used in each binding experiment. Optical densities of the control wells never exceeded 0.050 at all laminin concentrations employed for these experiments. The optical densities of the control wells were subtracted from the optical densities of the A β -containing wells that received similar laminin concentrations.

10 Non-specific absorbance obtained from A β containing wells that did not receive laminin were also subtracted from all data points. Thus, the equation in the form of: $OD_{exp} = OD_o + (S \times [laminin]) + (OD_{max} \times [laminin]/([laminin] + K_d))$ where $(S \times [laminin])$ represents non-specific binding (control wells) and OD_o is the non-specific absorbance, becomes $OD_{exp} = OD_{max} \times [laminin]/([laminin] + K_d)$. Therefore, at 50 % saturation $OD_{exp} = 0.50 OD_{max}$ and $K_d = [laminin]$. Determination of $[laminin]$ at 50% saturation was performed by non-linear

15 least square program (Ultrafit from Biosoft, UK) using a one-site model.

As demonstrated in Figure 1, EHS laminin bound immobilized A β (1-40) with a

20 single binding constant with an apparent dissociation constant of $K_d = 2.7 \times 10^{-9}$ M.

Several repeated experiments utilizing this solid phase binding immunoassay indicated that laminin bound A β (1-40) repetitively with one apparent binding constant.

Example 2

Inhibition of Alzheimer's Disease A β Fibril Formation by Laminin

5

The effects of laminin on A β fibrillogenesis was also determined using the previously described method of Thioflavin T fluorometry (Naiki et al, Lab. Invest. 65:104-110, 1991; Levine III, Protein Sci. 2:404-410, 1993; Levine III, Int. J. Exp. Clin. Invest. 2:1-6, 1995; Naiki and Nakakuki, Lab. Invest. 74:374-383, 1996). In this assay,

10

Thioflavin T binds specifically to fibrillar amyloid and this binding produces a fluorescence enhancement at 480 nm that is directly proportional to the amount of amyloid fibrils formed (Naiki et al, Lab. Invest. 65:104-110, 1991; Levine III, Protein Sci. 2:404-410, 1993;

15

Levine III, Int. J. Exp. Clin. Invest. 2:1-6, 1995; Naiki and Nakakuki, Lab. Invest. 74:374-383, 1996). In a first study, the effects of EHS laminin on A β (1-40) fibrillogenesis was assessed. For this study, 25 μ M of freshly solubilized A β (1-40)(Bachem Inc.,

Torrance, CA, USA; Lot # WM365) was incubated in microcentrifuge tubes at 37°C for 1 week (in triplicate), either alone, or in the presence of 100 nM EHS laminin (Sigma Chemical Company, St. Louis, MO, USA) in 100 mM Tris, 50 mM NaCl, pH 7.0 (TBS). 100 nM of laminin utilized for these studies represented a A β :laminin molar ratio of 250:1.

20

50 μ l aliquots were then taken from each tube for analysis at 1 hr, 1 day, 3 days, and 1 week. In a second set of studies, the effects of laminin on A β (1-40) fibril formation was directly compared to other basement membrane components including fibronectin, type IV collagen and perlecan. For these studies, 25 μ M of freshly solubilized A β (1-40) was

25

incubated in microcentrifuge tubes for 1 week (in triplicate) either alone, or in the presence of 100 nM of EHS perlecan (isolated as previously described)(Castillo et al, J. Biochem. 120:433-444, 1996), fibronectin (Sigma Chemical Company, St. Louis, MO, USA) or type

IV collagen (Sigma Chemical Company, St. Louis, MO, USA). 50 μ l aliquots were then taken for analysis at 1 hour, 1 day, 3 days and 1 week. In a third set of studies, 25 μ M of freshly solubilized A β (1-40) was incubated in microcentrifuge tubes for 1 week (in triplicate) either alone, or in the presence of increasing concentrations of laminin (i.e. 5 nM, 15 nM, 40 nM and 100 nM). 50 μ l aliquots were taken for analysis at 1 hour, 1 day, 3 days and 1 week.

For each determination described above, following each incubation period, A β peptides +/- laminin, perlecan, fibronectin or type IV collagen, were added to 1.2 ml of 100 μ M Thioflavin T (Sigma Chemical Co., St. Louis, MO) in 50 mM phosphate buffer (pH 6.0). Fluorescence emission at 480 nm was measured on a Turner instrument-model 450 fluorometer at an excitation wavelength of 450 nm. For each determination, the fluorometer was calibrated by zeroing in the presence of the Thioflavin T reagent alone, and by setting the 50 ng/ml riboflavin (Sigma Chemical Co., St. Louis, Mo) in the Thioflavin T reagent to 1800 fluorescence units. All fluorescence determinations were based on these references and any background fluorescence given off by laminin, perlecan, type IV collagen, or fibronectin alone in the presence of the Thioflavin T reagent was always subtracted from all pertinent readings.

As shown in Figure 2, freshly suspended A β (1-40) alone, following a 1 hour incubation at 37°C, demonstrated an initial fluorescence of 41 fluorescence units. During the 1 week incubation period there was a gradual increase in the fluorescence of 25 μ M A β (1-40) alone, increasing 6.7-fold from 1 hour to 1 week, with a peak fluorescence of 379 fluorescence units observed at 1 week. This increase was significantly inhibited when A β (1-40) was co-incubated with laminin, in comparison to A β alone. A β (1-40) co-incubated with laminin displayed fluorescence values that were 2.9-fold lower ($p < 0.001$) at 1 hour,

4.6-fold lower ($p<0.0001$) at 1 day, 30.6-fold lower ($p<0.0001$) at 3 days and 27.1-fold lower ($p<0.0001$) at 1 week. This study indicated that laminin was a potent inhibitor of A β amyloid fibril formation, nearly completely inhibiting amyloid fibril formation even after 1 week of incubation.

5

To determine whether the inhibitory effects of laminin was specific to this basement membrane component, an direct comparison was made to other known basement membrane components including perlecan, fibronectin, and type IV collagen. In these studies 25 μ M of A β (1-40) was incubated in the absence or presence of either 100 nM of laminin, 100 nM of fibronectin, 100 nM of type IV collagen and 100 nM of perlecan (Figure 3). Freshly solubilized A β (1-40) when incubated at 37°C gradually increased in fluorescence levels from 1 hour to 1 week (by 10.8-fold)(Figure 3), as previously demonstrated (Figure 2). Perlecan was found to significantly accelerate A β (1-40) amyloid formation at 1 day and 3 days, whereas fibronectin and type IV collagen only showed significant inhibition of A β (1-40) fibrillogenesis at 1 week. Laminin, on the other hand, was again found to be a very potent inhibitor of A β fibrillogenesis causing a 9-fold decrease at 1 and 3 days, and a 21-fold decrease at 1 week. This study reconfirmed the potent inhibitory effects of laminin on A β fibrillogenesis, and demonstrated the specificity of this inhibition, since none of the other basement membrane components (including fibronectin, type IV collagen and perlecan) were very effective inhibitors.

To determine whether the inhibitory effects of laminin on A β fibrillogenesis occurred in a dose-dependent manner, different concentrations of laminin (i.e. 5nM, 15 nM, 40 nM and 100 nM) were tested. As shown in Figure 4, freshly solubilized A β (1-40) when incubated at 37°C gradually increased from 1 hour to 1 week, as previously demonstrated (Figures 2 and 3). 100 nM of laminin significantly inhibited A β fibril formation at all time

points studied, including 1 hour, 1 day, 3 days and 7 days. Laminin was also found to inhibit A β fibril formation in a dose-dependent manner which was significant ($p < 0.05$) by 3 days of incubation. At 3 days and 7 days, both 100 nM and 40 nM of laminin significantly inhibited A β fibril formation. This study reconfirmed that laminin was a potent inhibitor of A β fibril formation and that this inhibition occurred in a dose-dependent manner.

Example 3

Laminin Causes Dose-Dependent Dissolution of Pre-Formed Alzheimer's Disease Amyloid Fibrils

The next study was implemented to determine whether laminin was capable of causing a dose-dependent dissolution of pre-formed Alzheimer's disease A β (1-40) amyloid fibrils. This type of activity would be important for any potential anti-Alzheimer's amyloid drug which can be used in patients who already have substantial amyloid deposition in brain. For example, Alzheimer's disease patients in mid-to late stage disease have abundant amyloid deposits in their brains as part of both neuritic plaques and cerebrovascular amyloid deposits. A therapeutic agent capable of causing dissolution of pre-existing amyloid would be advantageous for use in these patients who are at latter stages of the disease process.

For this study, 1 mg of A β (1-40)(Bachem Inc., Torrance, CA, USA; Lot #WM365) was dissolved in 1.0 ml of double distilled water (1mg/ml solution) and then incubated at 37°C for 1 week. 25 μ M of fibrillized A β was then incubated at 37°C in the presence or absence of laminin (from EHS tumor; Sigma Chemical Company, St. Louis, MO, USA) at concentrations of 125 nM, 63 nM, 31 nM and 16 nM containing 150 mM Tris HCl, 10 mM NaCl, pH 7.0. Following a 4 day incubation, 50 μ l aliquots were added

to 1.2ml of 100 μ M Thioflavin T (Sigma Chemical Co., St. Louis, MO) in 50mM NaPO₄ (pH 6.0) for fluorometry readings as described in example 2.

As shown in Figure 5, dissolution of pre-formed Alzheimer's disease A β amyloid fibrils by laminin occurred in a dose-dependent manner. A significant (p<0.001) 41% dissolution of pre-formed A β amyloid fibrils was observed with 125 nM of laminin, whereas 63 nM of laminin caused a significant (p<0.001) 39% dissolution. Furthermore, 31 nM and 16 nM of laminin still caused a significant (p<0.01) 28% and 25% dissolution of pre-formed A β amyloid fibrils. These data demonstrated that laminin causes dissolution of pre-formed Alzheimer's disease amyloid fibrils in a dose-dependent manner following a 4-day incubation.

Example 4

Laminin Does Not Significantly Inhibit Islet Amyloid Polypeptide (Amylin) Fibril Formation

In the next study, the specificity of the laminin inhibitory effects on Alzheimer's disease amyloid was determined by testing laminin's potential effects on another type of amyloid. Amyloid accumulation occurs in the islets of Langerhans in ~90% of patients with type II diabetes (Westermarck et al, Am. J. Path. 127:414-417, 1987). The major protein in islet amyloid is a 37 amino acid peptide, termed islet amyloid polypeptide or amylin which is known to be a normal secretory product of the beta-cells of the pancreas (Cooper et al, Proc. Natl. Acad. Sci., 84:8628-8632, 1987). The dose-dependent effects of laminin on amylin fibrillogenesis was determined using the Thioflavin T fluorometry assay. 25 μ M of A β (1-40)(Bachem Inc., Torrance, CA, USA; Lot #WM365) was incubated in microcentrifuge

tubes at 37°C for 1 week (in triplicate), either alone, or in the presence of 5 nM, 15 nM, 40 nM and 100 nM of laminin in 150 mM Tris HCl, 10 mM NaCl, pH 7.0 (TBS). 50 µl aliquots were taken from each tube for analysis at 1 hr, 1 day, 3 days, and 1 week using Thioflavin T fluorometry as described in example 2.

5

As shown in Figure 6, freshly suspended amylin alone following a 1-hour incubation at 37°C reached a maximum fluorescence of 1800 fluorescence units, which did not significantly change during the 1 week experimental period. The initial high fluorescence of amylin was attributed to amylin's ability to spontaneously form amyloid fibrils within a very short incubation period. Laminin at 100 nM did not significantly inhibit amylin fibril formation at all time points within the 1 week experimental period (Figure 6). In addition, no significant inhibition of amylin fibrillogenesis by laminin at decreasing concentrations (i.e. 40 nM, 15 nM and 5 nM) was observed, even though a decrease (but not significant) in amylin fibril formation was observed with 40 nM of laminin at 1 day, 3 days and 1 week (Figure 6). This study demonstrated that the inhibitory effects of laminin did not occur with amylin fibril formation, and demonstrated the specificity of the observed laminin inhibitory effects on Alzheimer's disease amyloid.

10

15

Example 5

20

Identification of V8 and Trypsin-Resistant Laminin Fragments which Interact with the Beta-Amyloid Protein of Alzheimer's Disease

25

In the next set of studies, we determined whether small fragment(s) of laminin generated by V8 or trypsin digestion would bind to Aβ. This would enable one to determine the domain(s) of laminin which bind Aβ and likely play a role in inhibition of Aβ fibril formation and causing dissolution of preformed Alzheimer's amyloid fibrils (as

demonstrated in the invention).

For these experiments, A β (1-40) was biotinylated according to the manufacturer's protocol (Pierce, Rockford, Illinois). For the ligand studies, intact EHS laminin was left undigested, or digested with V8 or trypsin (Sigma Chemical Company, St. Louis, MO, USA). More specifically, 2 μ g of trypsin or V8 protease in 2 μ l of 50 mM Tris-HCl buffer (pH 8.0) were added to 50 μ l of laminin (50 μ g)(in the same buffer) and incubated overnight at 37°C. The next day, 10 μ l of protease-digested laminin (or undigested laminin) was mixed with 10 μ l of 2X sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer, and heated for 5 minutes in a boiling water bath. SDS-PAGE was performed according to the method of Laemmli (Laemmli, U.K. Nature 227:680-685, 1970), or according to the method of Schagger and Jagow (Schagger and Jagow, Anal. Biochem. 166:368-379, 1987) using a Mini-Protean II electrophoresis system (Biorad) with precast 4-15% Tris-Glycine or 10-20% tricine polyacrylamide gels, respectively, and under non-reducing conditions. Electrophoresis occurred at 200V for 45 minutes along with pre-stained molecular weight standards.

After SDS-PAGE (10-20% tricine or 4-15% Tris-Glycine gels) was performed as described above, the separated laminin and its fragments (total protein of 10 μ g/lane) were transferred to polyvinylidene difluoride membrane (PVDF) using a Mini transblot electrophoresis transfer cell (Biorad, Hercules, CA, U.S.A.). Electrotransfer was performed at 100V for 2 hours. Following transfer, membranes were rinsed with methanol and dried. The fragment(s) of laminin involved in binding to A β were then detected by using biotinylated-A β (1-40), as described above. Blots were probed for 2 hours with 2 μ M biotinylated A β (1-40) in TTBS. The membranes were then rinsed three times (10 seconds each) with TTBS, probed for 30 minutes with streptavidin alkaline phosphatase conjugate

(Vectastain), rinsed again (as described above), and followed by the addition of an alkaline phosphatase substrate solution (Vectastain). Following color development, the reaction was stopped by flushing the membranes with double distilled water.

5 As shown in Figure 7, V8-digested laminin produced multiple protein fragments which interacted with biotinylated A β (1-40). Using a 4-15% Tris-Glycine gel system (Figure 7, lane 1), V8-resistant laminin fragments which interacted with A β included fragments of ~400 kDa (which probably represented intact laminin which was left undigested), ~100-130 kDa, ~85 kDa, and a prominent fragment at ~ 55 kDa. Using a 10-20% tricine gel system (Figure 7, lane 2), V8-resistant laminin fragments which interacted with A β included fragments of ~130 kDa, ~85 kDa, and a prominent fragment at ~ 55 kDa (Figure 7, lane 2, arrow). It is important to note that molecular size expressed in kilodaltons (kDa) are generally approximate. This study demonstrated that the smallest V8-resistant protein fragment of laminin which interacted with A β (1-40) was ~55 kDa.

15 As shown in Figure 8, trypsin-digested laminin produced multiple protein fragments which interacted with biotinylated A β (1-40). Using a 4-15% Tris-Glycine gel system (Figure 8, lane 1), trypsin-resistant laminin fragments which interacted with A β included fragments of ~400 kDa (which probably represented intact laminin which was left undigested) , ~150-200 kDa, ~97 kDa, ~65 kDa and a prominent fragment at ~ 30 kDa. Using a 10-20% tricine gel system (Figure 8, lane 2), trypsin-resistant laminin fragments which interacted with A β included fragments of ~97 kDa, ~90 kDa, ~65 kDa and a prominent fragment at ~ 30 kDa (Figure 8, lane 2, arrow). This study demonstrated that the smallest trypsin-resistant fragment of laminin which interacted with A β (1-40) was ~30 kDa.

Example 6

Identification of Elastase-Resistant Laminin Fragments Which Interact with the Beta-Amyloid Protein of Alzheimer's Disease

In the next set of studies, we determined whether small fragment(s) of laminin generated by elastase digestion would bind to A β . In addition, we sequenced and identified the region within elastase-resistant laminin which interacted with A β . For these experiments, A β (1-40) was biotinylated according to the manufacturer's protocol (Pierce, Rockford, Illinois). For the ligand studies, intact EHS laminin was left undigested, or digested with elastase (Sigma Chemical Company, St. Louis, MO, USA). For elastase digestion, 2 μ g of elastase in 8 μ l of 50 mM Tris-HCl buffer (pH 8.0) was added to 50 μ l of laminin (50 μ g)(in the same buffer) and incubated for 1.5 hours or 2.5 hours at 37°C. In addition, as a control, 2 μ g of elastase in 50 μ l of 50 mM Tris-HCl buffer (pH 8.0) was incubated for 2.5 hours at 37°C. Following the appropriate incubation times as described above, 10 μ l of each of the above incubations were mixed with 10 μ l of 2X SDS-PAGE electrophoresis sample buffer, and heated for 5 minutes in a boiling water bath. SDS-PAGE was performed according to the method of Laemmli (Laemmli, Nature 227:680-685, 1970) using a Mini-Protean II electrophoresis system with precast 4-15% Tris-Glycine polyacrylamide gels, and under non-reducing conditions. Electrophoresis occurred at 200V for 45 minutes along with pre-stained molecular weight standards (Biorad).

After SDS-PAGE was performed as described above, the separated laminin fragments were transferred to PVDF using a Mini transblot electrophoresis transfer cell (Millipore, Bedford, MA, U.S.A.). Electrotransfer was performed at 100V for 2 hours.

Following transfer, membranes were rinsed with methanol, dried and cut into two equal parts which were used for A β ligand blotting, or Coomassie blue staining and subsequent amino acid sequencing. The fragment(s) of laminin involved in binding to A β were then detected by using biotinylated-A β (1-40), as described above. Blots were probed for 2 hours with 2 μ M biotinylated A β (1-40) in TTBS. The membranes were then rinsed three times (10 seconds each) with TTBS, probed for 30 minutes with strepavidin alkaline phosphatase conjugate (Vectastain), rinsed again (as described above), and followed by the addition of an alkaline phosphatase substrate solution (Vectastain). Following color development, the reaction was stopped by flushing the membranes with double distilled water.

For Coomassie blue staining, PVDF membranes were immersed with 0.2% Coomassie Brilliant blue (w/v) in 50 % methanol, 10% acetic acid, and 40% distilled water for 2 minutes, and then rinsed with 50% methanol, 10% acetic acid, and 40% distilled water until visible bands were observed, and no background staining was present. The 55 kDa A β -binding laminin fragment, described below, was sent to the Biotechnology Service Center (Peptide Sequence Analysis Facility at the University of Toronto, Toronto, Ontario, Canada) and subjected to amino acid sequencing using a Porton 2090 Gas-Phase Microsequencer (Porton Instruments, Tarzana, CA) with on-line analysis of phenylthiohydantoin derivatives.

In Figure 9, Panel A represents an A β ligand blot whereas Panel B represents the equivalent Coomassie blue stained blot. As shown in Figure 9 (Panel A, lanes 2 and 3), elastase-digested laminin produced multiple protein fragments which bound biotinylated A β (1-40). Panel A, lane 1 represents undigested mouse EHS laminin, whereas lanes 2 and 3 represents laminin which had been digested with elastase for 1.5 hours or 2.5 hours,

respectively. Panel A, lane 4 represents elastase digestion for 2.5 hours in the absence of laminin. Undigested laminin (Fig. 9, Panel A, lane 1) which interacted with A β included multiple bands from > ~400 kDa to >~86 kDa, with the most prominent A β -interaction occurring with intact laminin (i.e. ~ 400 kDa). Elastase-resistant laminin protein fragments which interacted with A β (Fig. 9, Panel A, lanes 2 and 3) included fragments of >~400kDa, ~130 kDa (arrowhead), ~80-90 kDa, ~65 kDa and a prominent band at ~ 55 kDa (arrow). The interaction of these elastase-resistant laminin protein fragments with A β were only observed under non-reducing conditions suggesting that the A β interaction was also conformation dependent. The 130kDa elastase resistant laminin fragment which interacts with A β , is also believed to be part of the E8 fragment (see Figure 11), and is the same protein fragment of laminin that appears to be present in human serum and cerebrospinal fluid (see Examples 10 and 11). Figure 9, Panel A, lane 4 demonstrates that the band observed at ~29 kDa represents non-specific A β binding due to the presence of the elastase enzyme alone.

Figure 9, Panel B demonstrates all of the multiple protein bands which were stained by Coomassie blue. Note, for example, in Panel B, lanes 2 and 3, that elastase digestion of laminin produced multiple protein fragments between ~55 kDa and ~90 kDa which did not bind A β , and were not observed in the A β ligand blot (Fig. 9, Panel A, lanes 2 and 3).

Example 7

An A β -Binding Domain Within Laminin is Identified Within the Globular Repeats of the Laminin A Chain

The 55 kDa laminin fragment (ie. produced following 1.5 hours of elastase digestion) that demonstrated positive A β binding interaction by ligand blotting was then

prepared (Fig. 9, Panel B, lane 2, arrow) in large amounts for amino acid sequencing (as described in example 6). Sequence data determined the exact location within laminin that was involved in binding to A β . An 11-amino acid sequence was determined from sequencing of the 55 kDa band. The sequence identified was:

5 Leu-His-Arg-Glu-His-Gly-Glu-Leu-Pro-Pro-Glu (SEQ ID NO:1).

The specific A β -binding domain within laminin was then identified by comparison to known mouse laminin sequence (Sasaki and Yamada, J. Biol. Chem. 262:17111-17117, 1987; Sasaki et al, Proc. Natl. Acad. Sci. 84:935-939, 1987; Durkin, et al, Biochem. 27:5198-5204, 1988; Sasaki et al, J. Biol. Chem. 263:16536-16544, 1988), since mouse EHS laminin was utilized in the studies of the present invention. In addition, the complete amino acid sequence within laminin was retrieved from the National Center for Biotechnology Information, Bethesda, Maryland, U.S.A.

Figure 10 shows the complete amino acid sequence of mouse laminin A chain (Genebank accession number P19137; SEQ ID NO: 4). The 11 amino acid protein fragment sequenced from the ~55 kDa protein within laminin which binds A β is identified (Figure 10; bold underline and arrowhead; SEQ ID NO: 1) and matches exactly to the region within the third globular domain repeat of laminin A chain (Figure 11). The fourth globular domain repeat of mouse laminin A chain is shown as SEQ ID NO: 2 (Genebank Accession Number P19137; amino acids #2746-2922), whereas the fourth globular domain repeat of human laminin A chain is shown as SEQ ID NO: 3 (Genebank Accession Number P25391; amino acids #2737-2913).

Figure 11 shows two schematic representations of laminin (Colognato-Pyke et al, J. Biol. Chem. 270:9398-9406, 1995) and the newly discovered A β -binding region of laminin

(shown in left panel; between the two arrowheads) which is situated within the last three globular domains of the laminin A chain. The left panel of figure 11 illustrates laminin and fragments generated following protease digestions. Elastase fragments E1', E1X (dark line border), E-alpha-35 and E4 all correspond to regions of the short arms of laminin. Long arm fragments are E8, E3 and cathepsin G fragment C8-9. The E8 fragment produced by elastase digestion of laminin contains the long arm fragments containing the distal part of the long arm and the G subdomains 1-3, and consists of a 130-150 kDa (Yurchenco and Cheng, J. Biol. Chem. 268:17286-17299, 1993). The E3 fragments also produced by elastase digestion of laminin contains the distal long arm globule with G subdomains 4 and 5. The E3 fragment shown in Figure 11, Panel A, has previously shown to be a doublet at ~60 kDa and ~55 kDa (Yurchenco and Cheng, J. Biol. Chem. 268:17286-17299, 1993). This also confirms our discovery whereby the ~55 kDa fragment which we found to bind A β is localized within the E3 region of laminin (Figure 11, Left Panel).

The right panel of Figure 11 depicts the function map with the alpha (A chain), β (B1 chain) and gamma (B2 chain) chains of laminin shown in shades of decreasing darkness. EGF repeats are indicated by bars in the rod domains of the short arm. Domains, based on sequence analysis, are indicated in small Roman numerals and letters. The locations of heparin-binding, polymer-forming, and the active α 6 β 1 integrin-binding sites are shown in bold-face for the alpha-chain short arm. The long arm functions of heparin binding (heparin), α 6 β 1 integrin-recognition site (α 6 β 1), and dystroglycan (DG), mapped in other studies, are indicated in gray-shaded labels. It is interesting to note that the A β -binding region of laminin is also a region involved in binding to heparin.

It should also be emphasized that the globular domain repeats of the laminin A chain likely interacts with A β in a conformation dependent manner, since the interaction of the

~55-kilodalton elastase-resistant protein fragments with A β was only observed under non-reducing conditions.

Example 8

Identification of Laminin and Laminin Protein Fragments in Human Serum and Cerebrospinal Fluid Derived from Alzheimer's disease, Type II Diabetes, and/or Normal Aged Patients

In the next study, western blotting techniques using a polyclonal antibody against laminin was used to determine whether intact laminin and/or laminin fragments were present in human serum and cerebrospinal fluid obtained from Alzheimer's disease, type II diabetes and/or normal aged patients. In this study, human serum was obtained from the Alzheimer's disease Research Center at the University of Washington from either living aged patients who may have had corresponding mini-mental state examinations (where a score of 30 is normal, a score of 15 suggests moderate dementia and a score <10 suggests severe dementia), or from living aged patients who had subsequently died and were diagnosed at autopsy with Alzheimer's disease (following examination of their brains obtained postmortem). In addition, human serum was obtained from the Diabetes Endocrinology Research Center at the University of Washington. The following human serums were obtained and analyzed as part of this study: 1) patient #9; a normal 67 yr old female with a mini-mental score of 30; 2) patient #5226 - a 70 year old female with confirmed moderate Alzheimer's disease who also had a mini-mental score of 12 ; 3) patient #5211- a 66 year old male with confirmed Alzheimer's disease who also had a mini-mental score of 25; 4) patient B- a 63 year old male who had confirmed type II diabetes; 5) patient #5223- a 68 year old female with confirmed Alzheimer's disease who also had a mini-mental score of 22; 6) patient #22- an 83 yr old normal aged female who also had a mini-mental score of 30;

7) patient #C- a 68 year old male with confirmed type II diabetes. Each of these serums were utilized in this study and represent lanes 1-7 (left side) of Figure 12 (in the same order as above).

5 In addition, cerebrospinal fluid was obtained from the Alzheimer's disease Research Center at the University of Washington from either living aged patients who may have had corresponding mini-mental state examinations, or from living aged patients who had subsequently died and were diagnosed at autopsy with Alzheimer's disease (following examination of their brains obtained postmortem). The following human cerebrospinal
10 fluids were obtained as part of this study: 1) patient #6- a normal 64 year old female who had a mini-mental score of 30; 2) patient #7- a normal 67 year old male who had a mini-mental score of 30; 3) patient #8- a normal 80 year old female who had a mini-mental score of 30; 4) patient #9- a normal 67 year old female who had a mini-mental score of 30;
15 5) patient #1111P- a normal 78 year old female who had a mini-mental score of 30; 6) patient #50-a 66 year old male patient with probable moderate Alzheimer's disease as indicated by a mini-mental score of 15; 7) patient #54-a 73 year old male with probable severe Alzheimer's disease as indicated by a mini-mental score of 8. Each of these cerebrospinal fluid samples were utilized in this study and represent lanes 1-7 (right side) of Figure 12 (in the same order as above).

20 For the study described above, 10 μ l of human serum diluted at 1:10, or 10 μ l of undiluted human cerebrospinal fluid was added to 10 μ l of SDS-PAGE buffer and ligand blots were prepared as in Example 6. Blots were probed for 2 hours with a polyclonal antibody (used at a dilution of 1:10,000 in TTBS) against EHS laminin (Sigma Chemical
25 Company, St. Louis, MO). The membranes were then rinsed 3 times (10 seconds each) with TTBS and incubated for 1 hour with a biotinylated goat anti-rabbit IgG secondary

antibody diluted 1:1,000 with TTBS. The membranes were then rinsed three times (10 seconds each) with TTBS, probed for 30 minutes with strepavidin alkaline phosphatase conjugate (Vectastain), rinsed again (as described above), followed by the addition of an alkaline phosphatase substrate solution (Vectastain). Following color development, the reaction was stopped by flushing the membranes with double distilled water.

As shown in Figure 12, intact laminin (arrowheads) was present in human serum (lanes 1-7; left side) but not in human cerebrospinal fluid (lanes 1-7; right side). Qualitative observations suggest that intact laminin (as described above) may have been decreased in serum of Alzheimer's disease patients in comparison to controls (i.e. compare intact laminin in Figure 12, lane 1, left side-normal individual; to Figure 12, lane 2, left side-Alzheimer's disease patient). In addition to intact laminin, human serum derived from Alzheimer's disease, type II diabetes and normal aged patients also contained laminin immunoreactivity in a series of band from ~120 kDa to ~200 kDa (Figure 12, bands observed between the two arrows). On the other hand, cerebrospinal fluid samples did not contain intact laminin (Figure 12; lanes 1-7; right side) but only contained a series of laminin immunoreactive protein fragments from ~120 kDa to ~200 kDa (i.e. Figure 12, bands observed between the two arrows). This study determined that a series of laminin protein fragments are present in both human serum and cerebrospinal fluid of Alzheimer's disease, type II diabetes and normal aged patients, whereas intact laminin is only present in human serum. The novel discovery of the laminin fragments in human cerebrospinal fluid suggests that it may be used as a marker to determine the extent of laminin breakdown in the brain during Alzheimer's disease and other brain disorders.

Example 9

Identification of a ~130 Kilodalton Laminin Protein Fragment in Human Serum of Alzheimer's disease, Type II Diabetes and Normal Aged Patients which Binds A β

In the next study, A β ligand blotting techniques were utilized to identify whether laminin or laminin protein fragments present in human serum bind A β . In this study, human serum was obtained from the Alzheimer's disease Research Center at the University of Washington from either living patients who may have had corresponding mini-mental state examinations (where a score of 30 is normal, a score of 15 suggests moderate dementia and a score <10 suggests severe dementia), or from living patients who had subsequently died and were diagnosed at autopsy with Alzheimer's disease (following examination of their brains obtained postmortem). In addition, human serum was obtained from the Diabetes Endocrinology Research Center at the University of Washington. The first six human serum samples (i.e. Figure 13, lanes 1-6) were the same serum samples as indicated in Example 8. In addition, Figure 13 lanes 7-10 consisted of human serum obtained from lane 7) patient #E- a 54 year old male with confirmed type II diabetes, lane 8) patient #5230- a 72 year old female with confirmed moderate Alzheimer's disease who had a mini-mental score of 19, lane 9) patient #E-a 54 year old male with confirmed type II diabetes, and lane 10) patient #F- a 69 year old male with confirmed type II diabetes.

For this study, A β (1-40) was biotinylated according to the manufacturer's protocol (Pierce, Rockford, IL). For the ligand studies, following SDS-PAGE as described above in Example 8, separated laminin and its fragments present in human serum were transferred to polyvinylidene difluoride membrane (PVDF) using a Mini transblot electrophoresis transfer

cell. Electrotransfer was performed at 100V for 2 hours. Following transfer, membranes were rinsed with methanol and dried. The fragment(s) of laminin in human serum involved in binding to A β were then detected by using biotinylated-A β (1-40). Blots were probed for 2 hours with 1 μ M biotinylated A β (1-40) in TTBS. The membranes were then rinsed three times (10 seconds each) with TTBS, probed for 30 minutes with streptavidin alkaline phosphatase conjugate (Vectastain), rinsed again (as described above), and followed by the addition of an alkaline phosphatase substrate solution (Vectastain). Following color development, the reaction was stopped by flushing the membranes with double distilled water.

As shown in Figure 13, A β interacted with intact human laminin (arrow) in most samples of human serum. However, it was surprising to note that intact laminin was virtually absent in 2 of the 4 Alzheimer's disease patients serum (Fig. 13, lanes 5 and 8), suggesting that laminin-derived fragments may be important in Alzheimer's disease as a diagnostic marker. The most interesting discovery was that of all the laminin immunoreactive protein fragments found in human serum (i.e ~120 kDa to ~200 kDa, bands observed between the arrows, Figure 12, lanes 1-7, right side), only a prominent ~130 kDa band was found to interact with A β (Figure 13, arrowhead). This same prominent band is approximately the same molecular weight of the E8 band generated from mouse laminin following elastase digestion (see Figure 9), and which also contains the globular domain repeats of the laminin A chain. This study therefore determined that besides intact laminin, human serum contains a ~130 kDa laminin fragment which binds to A β , and may be important for keeping A β soluble in biological fluids such as blood. This study also suggests that qualitative and quantitative assessment of laminin fragments in human serum may prove diagnostic for the extent and progression of Alzheimer's disease, type II diabetes and other amyloidoses.

Example 10

Identification of a ~130 Kilodalton Laminin Protein Fragment in Human Cerebrospinal Fluid of Alzheimer's disease and Normal Aged Patients which Binds A β

In the next study, A β ligand blotting techniques were utilized to identify whether laminin protein fragments (<200 kDa) present in human cerebrospinal fluid bind A β . In this study, human cerebrospinal fluid was obtained from the Alzheimer's disease Research Center at the University of Washington from either living aged patients who may have had corresponding mini-mental state examinations (where a score of 30 is normal, a score of 15 suggests moderate Alzheimer's disease and a score <10 suggests moderate Alzheimer's disease), or from living aged patients who had subsequently died and were diagnosed at autopsy with Alzheimer's disease (following examination of their brains obtained postmortem). The following human cerebrospinal fluids were obtained and analyzed as part of this study (depicted in Figure 14, lanes 1-10): 1) patient #65- a 71 yr old male with probable severe Alzheimer's disease as indicated by a mini-mental score of 0; 2) patient #54- a 73 yr old male with probable severe Alzheimer's disease as indicated by a mini-mental score of 8.; 3) patient #6- a normal 64 yr old female who had a mini-mental score of 30; 4) patient #7- a normal 67 yr old male who had a mini-mental score of 30; 5) patient #8- a normal 80 yr old female who had a mini-mental score of 30; 6) patient #9- a normal 67 yr old female who had a mini-mental score of 30; 7) patient #1111P- a normal 78 yr old female who had a mini-mental score of 30; 8) patient #50-a 66 yr old male patient with probable moderate Alzheimer's disease as indicated by a mini-mental score of 15; 9) patient #52-a 69 yr old male with probable moderate Alzheimer's disease as indicated by a mini-mental score of 16; 10) patient #64-a 64 yr old male with probable severe Alzheimer's

disease as indicated by a mini-mental score of 0. Each of these cerebrospinal fluid samples were utilized in this study and represent lanes 1-10 of Figure 14 (in the same order as above).

5 For this study, A β ligand blotting was employed as described in Example 9. The fragment(s) of laminin in human cerebrospinal fluid involved in binding to A β were detected by using biotinylated-A β (1-40). Blots were probed for 2 hours with 50 nM of biotinylated A β (1-40) in TTBS. The rest of the A β ligand blotting procedure is as described above in Example 9.

10 As shown in Figure 14, A β interacted with laminin fragment bands between ~120 kDa and ~200 kDa in most samples of human cerebrospinal fluid. As observed in human serum, most samples of human cerebrospinal fluid also contained a prominent ~130 kDa laminin fragment (Figure 14, arrow) which interacted with A β . No intact A β -binding laminin was found in human cerebrospinal fluid (not shown), as previously demonstrated (Figure 12, Example 8). Again, this same prominent ~130 kDa A β -binding laminin fragment present in human cerebrospinal fluid is approximately the same molecular weight of the E8 band generated from laminin, and which also contains the globular domain repeats of the laminin A chain. This study therefore determined that human cerebrospinal fluid also contains a ~130 kDa laminin fragment which binds to A β , and may be important for keeping A β soluble in biological fluids such as cerebrospinal fluid.

Further Aspects and Utilizations of the Invention

Laminin-Derived Protein Fragments and Polypeptides

5 One therapeutic application of the present invention is to use laminin, laminin protein fragments which bind A β or other amyloid proteins, and/or laminin polypeptides derived from amino acid sequencing of the laminin fragments which bind A β (such as the ~130 kilodalton protein described herein) or other amyloid proteins, as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and
10 other amyloidoses. The amyloid diseases include, but are not limited to, the amyloid associated with Alzheimer's disease and Down's syndrome (wherein the specific amyloid is referred to as beta-amyloid protein or A β), the amyloid associated with chronic inflammation, various forms of malignancy and Familial Mediterranean Fever (wherein the specific amyloid is referred to as AA amyloid or inflammation-associated amyloidosis), the
15 amyloid associated with multiple myeloma and other B-cell dyscrasias (wherein the specific amyloid is referred to as AL amyloid), the amyloid associated with type II diabetes (wherein the specific amyloid is referred to as amylin or islet amyloid), the amyloid associated with the prion diseases including Creutzfeldt-Jakob disease, Gerstmann-
20 Straussler syndrome, kuru and animal scrapie (wherein the specific amyloid is referred to as PrP amyloid), the amyloid associated with long-term hemodialysis and carpal tunnel syndrome (wherein the specific amyloid is referred to as beta₂-microglobulin amyloid), the amyloid associated with senile cardiac amyloid and Familial Amyloidotic Polyneuropathy (wherein the specific amyloid is referred to as transthyretin or prealbumin), and the amyloid associated with endocrine tumors such as medullary carcinoma of the thyroid (wherein the
25 specific amyloid is referred to as variants of procalcitonin).

5 The polypeptides referred to above may be a natural polypeptide, a synthetic polypeptide or a recombinant polypeptide. The fragments, derivatives or analogs of the polypeptides to any laminin fragment referred to herein may be a) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue and such substituted amino acid residue may or may not be encoded by the genetic code, or b) one in which one or more of the amino acid residues includes a substituent group, or c) one in which the mature polypeptide is fused with another compound, such as a compound used to increase the half-life of the polypeptide (for example, polylysine), or d) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of the invention.

15 The tertiary structure of proteins refers to the overall 3-dimensional architecture of a polypeptide chain. The complexity of 3-dimensional structure arises from the intrinsic ability of single covalent bonds to be rotated. Rotation about several such bonds in a linear molecule will produce different nonsuperimposable 3-dimensional arrangements of the atoms that are generally described as conformations.

20 Protein conformation is an essential component of protein-protein, protein-substrate, protein-agonist, protein-antagonist interactions. Changes in the component amino acids of protein sequences can result in changes that have little or no effect on the resultant protein conformation. Conversely, changes in the peptide sequences can have effects on the protein conformation resulting in reduced or increased protein-protein, etc. interactions. Such changes and their effects are generally disclosed in Proteins: Structures and Molecular Properties by Thomas Creighton W.H. Freeman and Company, New York, 1984 which

is hereby incorporated by reference.

“Conformation” and “conformation similarity” when used in this specification and claims refers to a polypeptide’s ability (or any other organic or inorganic molecule) to assume a given shape, through folding and the like, so that the shape, or conformation, of the molecule becomes an essential part of it’s functionality, sometimes to the exclusion of its chemical makeup. It is generally known that in biological processes two conformational similar molecules may be interchangeable in the process, even the chemically different. “Conformational similarity” refers to the latter interchangeability or substitutability. For example, laminin and laminin-derived protein fragments are among the subjects of the invention because they have been shown to bind the A β protein and render it inactive in fibril formation; it is contemplated that other molecules that are conformationally similar to laminin, or any claimed laminin fragment or polypeptide, may be substituted in the claimed method to similarly render the A β inactive in fibrillogenesis and other amyloid processes. In general it is contemplated that levels of conformational similarity at or above 70% are sufficient to assume homologous functionality in the claimed processes, though reduced levels of conformational similarity may be made to serve as well. Conformational similar levels at or above 90% should provide some level of additional homologue functionality.

Thus, one skilled in the art would envisage that changes can be made to the laminin sequence, or fragments or polypeptides thereof, that would increase, decrease or have no effect on the binding of laminin or fragments thereof, to A β amyloid. In addition, one skilled in the art would envisage various post-translational modifications such as phosphorylation, glycosylation and the like would alter the binding of laminin, laminin fragments or laminin polypeptides to A β amyloid.

The polypeptides of the present invention include the polypeptides or fragments of laminin described herein, including but not limited to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and fragments thereof, as well as polypeptides which have at least 70% similarity (preferably 70 % identity) and more preferably a 90% similarity (more preferably a 90% identity) to the polypeptides described above.

Fragments or portions of the polypeptides or fragments of laminin of the present invention may be employed for producing the corresponding full-length polypeptides by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full length polypeptides.

The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue.

Chemical polypeptide synthesis is a rapidly evolving area in the art, and methods of solid phase polypeptide synthesis are well-described in the following references, hereby entirely incorporated by reference (Merrifield, J. Amer. Chem. Soc. 85:2149-2154, 1963; Merrifield, Science 232:341-347, 1986; Fields, Int. J. Polypeptide Prot. Res. 35, 161, 1990).

Recombinant production of laminin polypeptides can be accomplished according to known method steps. Standard reference works setting forth the general principles of recombinant DNA technology include Watson, Molecular Biology of the Gene, Volumes I and II, The Benjamin/Cummings Publishing Company Inc., publisher, Menlo Park, Calif. 1987; Ausubel et al, eds., Current Protocols in Molecular Biology, Wiley Interscience, publisher, New York, N.Y. 1987; 1992; and Sambrook et al, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory, publisher, Cold Spring Harbor, N.Y. 1989, the entire contents of which references are herein incorporated by reference.

The polypeptides of the present invention may also be utilized as research reagents and materials for discovery of treatments and diagnostics for human diseases.

Antibodies

Antibodies generated against the polypeptides corresponding to specific sequences recognizing the laminin fragments of the present invention which bind A β or other amyloid proteins can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptides from tissue expressing that polypeptide. Preferred embodiments include, but are not limited to, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, and

fragments thereof, as well as polypeptides which have at least 70% similarity (preferably 70 % identity) and more preferably a 90% similarity (more preferably a 90% identity) to the polypeptides described above.

5 The term "antibody" is meant to include polyclonal antibodies, monoclonal antibodies, chimeric antibodies, anti-idiotypic antibodies to antibodies specific for laminin-derived protein fragments or polypeptides of the present invention.

10 Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen.

15 A monoclonal antibody contains a substantially homogeneous population of antibodies specific to antigens, which population contains substantially similar epitope binding sites. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, Nature 256:495-497, 1975), the trioma technique, the human B-cell hybridoma technique (Kozbor et al, Immunology Today 4:72, 1983), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp.77-96, 1985).

20 Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, GILD and any subclass thereof.

25 Chimeric antibodies are molecules different portions of which are derived from different animal species, such as those having variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region, which are primarily used to reduce immunogenicity in application and to increase yields in production. Chimeric

antibodies and methods for their production are known in the art (ex. Cabilly et al, Proc. Natl. Acad. Sci. U.S.A. 81:3273-3277, 1984; Harlow and Lane: Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory 1988).

5 An anti-idiotypic antibody is an antibody which recognizes unique determinants generally associated with the antigen-binding site of an antibody. An anti-idiotypic antibody can be prepared by immunizing an animal of the same species and genetic type (e.g., mouse strain) as the source of the monoclonal antibody with the monoclonal antibody to which an anti-idiotypic antibody is being prepared. The immunized animal will
10 recognize and respond to the idiotypic determinants of the immunizing antibody by producing an antibody to these idiotypic determinants (the anti-idiotypic antibody). See, for example, U.S. Patent No. 4,699,880, which is herein incorporated by reference.

15 The term "antibody" is also meant to include both intact molecules as well as fragments thereof, such as, for example, Fab and F(ab')₂, which are capable of binding antigen. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody (Wahl et al, J. Nucl. Med. 24:316-325, 1983).

20 The antibodies or fragments of antibodies, useful in the present invention may be used to quantitatively or qualitatively detect laminin or laminin-derived fragments in a sample or to detect presence of cells which express a laminin polypeptide of the present invention. This can be accomplished by immunofluorescence techniques employing a
25 fluorescently labeled antibody coupled with light microscopic, flow cytometric or fluorometric detection.

One of the ways in which a laminin fragment antibody can be detectably labeled is by linking the same to an enzyme and use in an enzyme immunoassay (EIA). This enzyme, in turn, when later exposed to an appropriate substrate, will react with the substrate in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorometric, or by visual means. Enzymes which can be used detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colometric methods which employ a chromogenic substrate for the enzyme. Detection can be accomplished by colometric methods which employ a chromogenic substrate for the enzyme. Detection can also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate with similarly prepared standards (see Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory 1988; Ausubel et al, eds., Current Protocols in Molecular Biology, Wiley Interscience, N.Y. 1987, 1992).

Detection may be accomplished using any of a variety of other immunoassays. For example, by radiolabeling of the antibodies or antibody fragments, it is possible to detect R-PTPase through the use of a radioimmunoassay (RIA). A good description of RIA may be found in Laboratory Techniques and Biochemistry in Molecular Biology, by Work et al, North Holland Publishing Company, NY (1978) with particular reference to the chapter entitled "An Introduction to Radioimmune Assay and Related Techniques" by Chard, incorporated entirely by reference herein. The radioactive isotope can be detected by such means as the use of a gamma-counter, a scintillation counter or by autoradiography.

It is also possible to label a laminin fragment polypeptide antibody with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labelling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine, commercially available, e.g., from Molecular Probes, Inc. (Eugene, Oregon, U.S.A.).

The antibody can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or other of the lanthanide series. These metals can be attached to the antibody using such metal groups as diethylenetriamine pentaacetic acid (EDTA).

The antibody can also be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, therrromatic acridinium ester, imidazole, acridinium salt, and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

The antibodies (or fragments thereof) useful in the present invention may be

employed histologically, as in immunofluorescence or immunoelectron microscopy, for in situ detection of a laminin fragment of the present invention. In situ detection may be accomplished by removing a histological specimen from a patient, and providing the labeled antibody of the present invention to such a specimen. The antibody (or fragment) is preferably provided by applying or by overlaying the labeled antibody (or fragment) to a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of a laminin fragment polypeptide but also its distribution on the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

In accordance with yet a further aspect of the present invention there are provided antibodies against laminin, laminin fragments and/or laminin-derived polypeptides which interact with A β or other amyloid proteins, or derivatives thereof. These antibodies can be used for a number of important diagnostic and/or therapeutic applications as described herein. In one aspect of the invention, polyclonal and/or monoclonal antibodies made against laminin, laminin fragments and/or laminin-derived polypeptides which bind A β or other amyloid proteins, may be utilized for Western blot analysis (using standard Western blotting techniques knowledgeable to those skilled in the art) to detect the presence of amyloid protein-binding laminin fragments or amyloid protein-binding laminin polypeptides in human tissues and in tissues of other species. Western blot analysis can also be used to determine the apparent size of each amyloid protein-binding laminin fragment. In addition, Western blotting following by scanning densitometry (known to those skilled in the art) can be used to quantitate and compare levels of each of the laminin fragments in tissue samples, biological fluids or biopsies obtained from individuals with specific diseases (such as the amyloid diseases) in comparison to tissue samples, biological fluids or biopsies obtained

from normal individuals or controls. Biological fluids, include, but are not limited to, blood, plasma, serum, cerebrospinal fluid, sputum, saliva, urine and stool.

In yet another aspect of the invention, polyclonal and/or monoclonal antibodies made against laminin, laminin fragments and/or laminin-derived peptides which bind A β or other amyloid proteins, can be utilized for immunoprecipitation studies (using standard immunoprecipitation techniques known to one skilled in the art) to detect laminin, laminin fragments and/or laminin-derived peptides which bind A β or other amyloid proteins, in tissues, cells and/or biological fluids. Use of the laminin, laminin fragment and/or laminin-derived peptide antibodies for immunoprecipitation studies can also be quantitated to determine relative levels of laminin, laminin fragments and/or laminin-derived peptides which interact with A β or other amyloid proteins, in tissues, cells and/or biological fluids. Quantitative immunoprecipitation can be used to compare levels of laminin, laminin fragments and/or laminin amyloid protein-binding peptides in tissue samples, biological fluids or biopsies obtained from individuals with specific diseases (such as the amyloid diseases) in comparison to tissue samples, biological fluids or biopsies obtained from normal individuals or controls.

Therapeutic Applications

Yet another aspect of the present invention is to make use of laminin, laminin fragments and/or laminin-derived polypeptides as amyloid inhibitory therapeutic agents. The laminin-derived peptide sequences or fragments can be synthesized utilizing standard techniques (ie. using an automated synthesizer). Laminin, laminin fragments and/or laminin-derived polypeptides which bind A β or other amyloid proteins, can be used as potential blocking therapeutics for the interaction of laminin in a number of biological processes and

diseases (such as in the amyloid diseases described above). In a preferred embodiment, specific peptides made against the amino acid sequence of laminin contained within the ~55 kDa laminin fragment (i.e. globular repeats within the laminin A chain; SEQ ID NO 3) described in the present invention, may be used to aid in the inhibition of amyloid formation, deposition, accumulation, and /or persistence in a given patient. Likewise, in another preferred embodiment anti-idiotypic antibodies made against laminin, laminin fragments and/or laminin-derived polypeptides (as described above) may be given to a human patient as potential blocking antibodies to disrupt continued amyloid formation, deposition, accumulation and/or persistence in the given patient.

Preparations of laminin-derived polypeptides for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions, which may contain axillary agents or excipients which are known in the art. Pharmaceutical compositions such as tablets, pills, tablets, caplets, soft and hard gelatin capsules, lozenges, sachets, cachets, vegicaps, liquid drops, elixers, suspensions, emulsions, solutions, syrups, tea bags, aerosols (as a solid or in a liquid medium), suppositories, sterile injectable solutions, sterile packaged powders, can be prepared according to routine methods and are known in the art.

In yet another aspect of the invention, laminin, laminin fragments and/or laminin-derived polypeptides may be used as an effective therapy to block amyloid formation, deposition, accumulation and/or persistence as observed in the amyloid diseases. For example, the invention includes a pharmaceutical composition for use in the treatment of amyloidoses comprising a pharmaceutically effective amount of a laminin, laminin fragment and/or laminin-derived polypeptide anti-idiotypic antibody and a pharmaceutically acceptable carrier. The compositions may contain the laminin, laminin fragments and/or laminin-derived polypeptide anti-idiotypic antibody, either unmodified, conjugated to a potentially

therapeutic compound, conjugated to a second protein or protein portion or in a recombinant form (ie. chimeric or bispecific laminin, laminin fragment and/or laminin polypeptide antibody). The compositions may additionally include other antibodies or conjugates. The antibody compositions of the invention can be administered using conventional modes of administration including, but not limited to, topical, intravenous, intra-arterial, intraperitoneal, oral, intralymphatic, intramuscular or intralumbar. Intravenous administration is preferred. The compositions of the invention can be a variety of dosage forms, with the preferred form depending upon the mode of administration and the therapeutic application. Optimal dosage and modes of administration for an individual patient can readily be determined by conventional protocols.

Laminin, laminin-derived protein fragments, and laminin-derived polypeptides, or antibodies of the present invention may be administered by any means that achieve their intended purpose, for example, to treat laminin involved pathologies, such as Alzheimer's disease and other amyloid diseases, or other related pathologies, using a laminin-derived polypeptide described herein, in the form of a pharmaceutical composition.

For example, administration of such a composition may be by various parenteral routes such as subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intranasal, transdermal or buccal routes. Alternatively, or concurrently, administration may be by the oral route. Parenteral administration can be by bolus injection or by gradual perfusion over time.

A preferred mode of using a laminin-derived polypeptide, or antibody pharmaceutical composition of the present invention is by oral administration or intravenous application.

A typical regimen for preventing, suppressing or treating laminin-involved pathologies, such as Alzheimer's disease amyloidosis, comprises administration of an effective amount of laminin-derived polypeptides, administered over a period of one or several days, up to and including between one week and about 24 months.

It is understood that the dosage of the laminin-derived polypeptides of the present invention administered in vivo or in vitro will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. The most preferred dosage will be tailored to the individual subject, as is understood and determinable by one of skill in the art, without undue experimentation.

The total dose required for each treatment may be administered by multiple doses or in a single dose. A laminin-derived polypeptide may be administered alone or in conjunction with other therapeutics directed to laminin-involved pathologies, such as Alzheimer's disease or amyloid diseases, as described herein.

Effective amounts of a laminin-derived polypeptide or composition, which may also include a laminin-fragment derived antibody, are about 0.01 μ g to about 100mg/kg body weight, and preferably from about 10 μ g to about 50 mg/kg body weight, such as 0.05, 0.07, 0.09, 0.1, 0.5, 0.7, 0.9., 1, 2, 5, 10, 20, 25, 30, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 mg/kg.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions, which may contain axillary agents or excipients

which are known in the art. Pharmaceutical compositions comprising at least one laminin-derived polypeptide, such as 1-10 or 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 laminin-derived polypeptides, of the present invention may include all compositions wherein the laminin-derived polypeptide is contained in an amount effective to achieve its intended purpose. In addition to at least one laminin-derived polypeptide, a pharmaceutical composition may contain suitable pharmaceutically acceptable carriers, such as excipients, carriers and/or axillaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically.

Pharmaceutical compositions comprising at least one laminin-derived polypeptide or antibody may also include suitable solutions for administration intravenously, subcutaneously, dermally, orally, mucosally, rectally or may by injection or orally, and contain from about 0.01 to 99 percent, preferably about 20 to 75 percent of active component (i.e. polypeptide or antibody) together with the excipient. Pharmaceutical compositions for oral administration include pills, tablets, caplets, soft and hard gelatin capsules, lozenges, sachets, cachets, vegicaps, liquid drops, elixers, suspensions, emulsions, solutions, and syrups.

The laminin, laminin-derived protein fragments, and laminin-derived polypeptides for Alzheimer's disease and other central nervous system amyloidoses may be optimized to cross the blood-brain barrier. Methods of introductions include but are not limited to systemic administration, parenteral administration i.e., via an intraperitoneal, intravenous, perioral, subcutaneous, intramuscular, intraarterial, intradermal, intramuscular, intranasal, epidural and oral routes. In a preferred embodiment, laminin, laminin-derived protein fragments, and laminin-derived polypeptides may be directly administered to the cerebrospinal fluid by intraventricular injection. In a specific embodiment, it may be

desirable to administer laminin, laminin-derived protein fragments, and laminin-derived polypeptides locally to the area or tissue in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, by injection, by infusion using a cannulae with osmotic pump, by means of a catheter, by means of a suppository, or by means of an implant.

In yet another embodiment laminin, laminin-derived protein fragments, and laminin-derived polypeptides may be delivered in a controlled release system, such as an osmotic pump. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, ie. the brain, thus requiring only a fraction of the systemic dose.

In yet another aspect of the present invention, peptidomimetic compounds modelled from laminin, laminin fragments and/or laminin-derived polypeptides identified as binding A β or other amyloid proteins, may serve as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses. Peptidomimetic modelling is implemented by standard procedures known to those skilled in the art.

In yet another aspect of the present invention, compounds that mimic the 3-dimensional A β binding site on laminin using computer modelling, may serve as potent inhibitors of amyloid formation, deposition, accumulation and/or persistence in Alzheimer's disease and other amyloidoses. Design and production of such compounds using computer modelling technologies is implemented by standard procedures known to those skilled in the art.

Recombinant DNA technology, including human gene therapy, has direct

applicability to the laminin proteins and their fragments, of this invention. One skilled in the art can take the peptide sequences disclosed herein and create corresponding nucleotide sequences that code for the corresponding peptide sequences. These sequences can be cloned into vectors such as retroviral vectors, and the like. These vectors can, in turn, be transfected into human cells such as hepatocytes or fibroblasts, and the like. Such transfected cells can be introduced into humans to treat amyloid diseases. Alternatively, the genes can be introduced into the patients directly. The basic techniques of recombinant DNA technology are known to those of ordinary skill in the art and are disclosed in Recombinant DNA Second Edition, Watson, et al., W.H. Freeman and Company, New York, 1992, which is hereby incorporated by reference.

Diagnostic Applications

Another aspect of the invention is to provide polyclonal and/or monoclonal antibodies against laminin, laminin fragments and/or laminin-derived polypeptides which bind A β or other amyloid proteins, which would be utilized to specifically detect laminin, laminin fragments and/or laminin-derived peptides in human tissues and/or biological fluids. In one preferred embodiment, polyclonal or monoclonal antibodies made against a peptide portion or fragment of laminin, can be used to detect and quantify laminin, laminin fragments and/or laminin-derived polypeptides in human tissues and/or biological fluids. Polyclonal and/or monoclonal peptide antibodies can also be utilized to specifically detect laminin fragments and/or laminin-derived polypeptides in human tissues and/or biological fluids. In a preferred embodiment, a polyclonal or monoclonal antibody made specifically against a peptide portion or fragment of ~55 kDa elastase-resistant protein which binds A β (as described herein), can be used to detect and quantify this laminin fragment in human tissues and/or biological fluids. In another preferred embodiment, a polyclonal or

monoclonal antibody made specifically against a peptide portion or fragment of ~130 kDa laminin-derived protein which is present in human biological fluids and binds A β (as described herein), can be used to detect and quantify this laminin fragment in human tissues and/or biological fluids. Other preferred embodiments include, but are not limited to, making polyclonal or monoclonal antibodies made specifically against a peptide portion or fragment of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, as well as polypeptides which have at least 70% similarity (preferably 70 % identity) and more preferably a 90% similarity (more preferably a 90% identity) to the polypeptides described above.

For detection of laminin fragments and/or laminin-derived polypeptides described above in human tissues, cells, and/or in cell culture, the polyclonal and/or monoclonal antibodies can be utilized using standard immunohistochemical and immunocytochemical techniques, known to one skilled in the art.

For detection and quantitation of laminin, laminin fragments and/or laminin-derived polypeptides in biological fluids, including cerebrospinal fluid, blood, plasma, serum, urine, sputum, and/or stool, various types of ELISA assays can be utilized, known to one skilled in the art. An antibody molecule of the present invention may be adapted for utilization in an immunometric assay, also known as a "two-site" or "sandwich" assay. In a typical immunometric assay, a quantity of unlabeled antibody (or fragment of antibody) is bound to a solid support or carrier, and a quantity of detectable labeled soluble antibody is added to permit detection and/or quantitation of the ternary complex formed between solid-phase antibody, antigen, and labeled antibody.

In a preferred embodiment, a "sandwich" type of ELISA can be used. Using this preferred method a pilot study is first implemented to determine the quantity of binding of each laminin-fragment monoclonal antibody to microtiter wells. Once this is determined, aliquots (usually in 40 μ l of TBS; pH 7.4) of the specific laminin-fragment antibody are allowed to bind overnight to microtiter wells (Maxisorb C plate from Nunc) at 4°C. A series of blank wells not containing any laminin-fragment specific monoclonal antibody are also utilized as controls. The next day, non-bound monoclonal antibody is shaken off the microtiter wells. All of the microtiter wells (including the blank wells) are then blocked by incubating for 2 hours with 300 μ l of Tris-buffered saline containing 0.05% Tween-20 (TTBS) plus 2% bovine serum albumin, followed by 5 rinses with TTBS. 200 μ l of cerebrospinal fluid, blood, plasma, serum, urine, sputum, and/or stool and/or any other type of biological sample is then diluted (to be determined empirically) in TTBS containing 2% bovine serum albumin and placed in wells (in triplicate) containing bound laminin-fragment antibody (or blank) and incubated for 2 hours at room temperature. The wells are then washed 5 times with TTBS. A second biotinylated-monoclonal antibody against the same laminin-derived fragment (but which is against a different epitope) is then added to each well (usually in 40 μ l of TBS; pH 7.4) and allowed to bind for 2 hours at room temperature to any laminin-fragment captured by the first antibody. Following incubation, the wells are washed 5 times with TTBS. Bound materials are then detected by incubating with 100 μ l of peroxidase-avidin complex (1:250 dilution in TTBS with 0.1% BSA) for 1 hour on a rotary shaker. After 5 washes with TTBS, a substrate solution (100 μ l, OPD-Sigma Fast from Sigma Chemical Co., St. Louis, MO, USA) is added and allowed to develop significant color (usually 8-10 minutes). The reaction is stopped with 50 μ l of 4N sulfuric acid and read on a standard spectrophotometer at 490 nm. This ELISA can be utilized to determine differences in specific laminin fragments (and/or A β -binding laminin fragments) in biological fluids which can serve as a diagnostic marker to follow the

progression on a live patient during the progression of disease (ie. monitoring of amyloid disease as an example). In addition, quantitative changes in laminin fragments can also serve as a prognostic indicator monitoring how a live patient will respond to treatment which targets a given amyloid disease such as Alzheimer's disease. Such assays can be provided in a kit form.

A competition assay may also be employed wherein antibodies specific to laminin, laminin fragments and/or laminin-derived polypeptides are attached to a solid support and labelled laminin, laminin fragments and/or laminin-derived polypeptides and a sample derived from a host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to the quantity of laminin, laminin fragments and/or laminin-derived polypeptides in the sample. This standard technique is known to one skilled in the art.

Another object of the present invention is to use laminin, laminin fragments and/or laminin-derived polypeptides, in conjunction with laminin, laminin fragment and/or laminin-derived peptide antibodies, in an ELISA assay to detect potential laminin, laminin fragment and/or laminin-derived peptide autoantibodies in human biological fluids. Such a diagnostic assay may be produced in a kit form. In a preferred embodiment, peptides containing the sequences of laminin, laminin-derived fragments and laminin-derived polypeptides as in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11, as well as polypeptides which have at least 70% similarity (preferably 70 % identity) and more preferably a 90% similarity (more preferably a 90% identity) to the polypeptides described above, will be used to initially bind to microtiter wells in an ELISA plate. A pilot study is first implemented to determine the quantity of binding of each laminin fragment

polypeptide to microtiter wells. Once this is determined, aliquots (usually 1-2µg in 40 µl of TBS; pH 7.4) of specific laminin fragment polypeptides (as described herein) are allowed to bind overnight to microtiter wells (Maxisorb C plate from Nunc) at 4°C. All the microtiter wells (including blank wells without the laminin fragment polypeptides) are blocked by incubating for 2 hours with 300 µl of Tris-buffered saline (pH 7.4) with 0.05% Tween-20 (TTBS), containing 2% albumin. This is followed by 5 rinses with TTBS. The patients' biological fluids (i.e., cerebrospinal fluid, blood, plasma, serum, sputum, urine, and/or stool) are then utilized and 200 µl are diluted (to be determined empirically) with TTBS containing 2% bovine serum albumin, and placed in microtiter wells (in triplicate) containing a specific laminin fragment polypeptide or blank wells (which do not contain peptide), and are incubated at 1.5 hours at room temperature. Any autoantibodies present in the biological fluids against the laminin fragment will bind to the substrate bound laminin fragment polypeptide (or fragments thereof). The wells are then rinsed by washing 5 times with TTBS. 100 µl of biotinylated polyclonal goat anti-human IgGs (Sigma Chemical company, St. Louis, MO, USA), diluted 1:500 in TTBS with 0.1% bovine serum albumin, is then aliquoted into each well. Bound materials are detected by incubating with 100 µl of peroxidase-avidin complex (1:250 dilution in TTBS with 0.1% bovine serum albumin) for 1 hour on a rotary shaker. Following 5 washes with TTBS, substrate solution (100 µl, OPD-Sigma Fast from Sigma Chemical Company, St. Louis, MO, USA) is added and allowed to develop significant color (usually 8-10 minutes). The reaction is stopped with 50 µl of 4N sulfuric acid added to each well and read on a standard spectrophotometer at 490 nm. This assay system can be utilized to not only detect the presence of autoantibodies against laminin fragments in biological fluids, but also to monitor the progression of disease by following elevation or diminution of laminin fragment autoantibody levels. It is believed that patients demonstrating excessive laminin fragment formation, deposition, accumulation and/or persistence as may be observed in the amyloid diseases, will also carry autoantibodies

against the laminin fragments in their biological fluids. Various ELISA assay systems, knowledgeable to those skilled in the art, can be used to accurately monitor the degree of laminin fragments in biological fluids as a potential diagnostic indicator and prognostic marker for patients during the progression of disease (ie. monitoring of an amyloid disease for example). Such assays can be provided in a kit form. In addition, quantitative changes in laminin fragment autoantibody levels can also serve as a prognostic indicator monitoring how a live patient will respond to treatment which targets a given amyloid disease.

Other diagnostic methods utilizing the invention include diagnostic assays for measuring altered levels of laminin, laminin fragments and/or laminin-derived polypeptides in various tissues compared to normal control tissue samples. Assays used to detect levels of laminin, laminin fragments and/or laminin-derived polypeptides in a sample derived from a host are well-known to those skilled in the art and included radioimmunoassays, competitive-binding assays, Western blot analysis and preferably ELISA assays (as described above).

Yet another aspect of the present invention is to use the antibodies recognizing laminin, laminin fragments and/or laminin-derived polypeptides for labellings, for example, with a radionucleotide, for radioimaging or radioguided surgery, for in vivo diagnosis, and/or for in vitro diagnosis. In one preferred embodiment, radiolabelled peptides or antibodies made (by one skilled in the art) against laminin, laminin fragments and/or laminin-derived polypeptides may be used as minimally invasive techniques to locate laminin, laminin fragments and/or laminin-derived polypeptides, and concurrent amyloid deposits in a living patient. These same imaging techniques could then be used at regular intervals (ie. every 6 months) to monitor the progression of the amyloid disease by following the specific levels of laminin, laminin fragments and/or laminin-derived

polypeptides.

Yet another aspect of the present invention is to provide a method which can evaluate a compound's ability to alter (diminish or eliminate) the affinity of a given amyloid protein (as described herein) or amyloid precursor protein, to laminin, laminin-derived fragments or laminin-derived polypeptides. By providing a method of identifying compounds which affect the binding of amyloid proteins, or amyloid precursor proteins to such laminin-derived fragments, the present invention is also useful in identifying compounds which can prevent or impair such binding interaction. Thus, compounds can be identified which specifically affect an event linked with the amyloid formation, amyloid deposition, and/or amyloid persistence condition associated with Alzheimer's disease and other amyloid diseases as described herein.

According to one aspect of the invention, to identify for compounds which allow the interaction of amyloid proteins or precursor proteins to laminin-derived fragments or laminin polypeptides, either amyloid or laminin fragments are immobilized, and the other of the two is maintained as a free entity. The free entity is contacted with the immobilized entity in the presence of a test compound for a period of time sufficient to allow binding of the free entity to the immobilized entity, after which the unbound free entity is removed. Using antibodies which recognize the free entity, or other means to detect the presence of bound components, the amount of free entity bound to immobilized entity can be measured. By performing this assay in the presence of a series of known concentrations of test compound and, as a control, the complete absence of test compound, the effectiveness of the test compound to allow binding of free entity to immobilized entity can be determined and a quantitative determination of the effect of the test compound on the affinity of free entity to immobilized entity can be made. By comparing the binding affinity of the amyloid-laminin fragment

complex in the presence of a test compound to the binding affinity of the amyloid-laminin fragment complex in the absence of a test compound, the ability of the test compound to modulate the binding can be determined.

5 In the case in which the amyloid is immobilized, it is contacted with free laminin-derived fragments or polypeptides, in the presence of a series of concentrations of test compound. As a control, immobilized amyloid is contacted with free laminin-derived polypeptides, or fragments thereof in the absence of the test compound. Using a series of concentrations of laminin-derived polypeptides, the dissociation constant (K_d) or other
10 indicators of binding affinity of amyloid-laminin fragment binding can be determined. In the assay, after the laminin-derived polypeptides or fragments thereof is placed in contact with the immobilized amyloid for a sufficient time to allow binding, the unbound laminin polypeptides are removed. Subsequently, the level of laminin fragment-amyloid binding can be observed. One method uses laminin-derived fragment antibodies, as described in the invention, to detect the amount of specific laminin fragments bound to the amyloid or the amount of free laminin fragments remaining in solution. This information is used to
15 determine first qualitatively whether or not the test compound can allow continued binding between laminin-derived fragments and amyloid. Secondly, the data collected from assays performed using a series of test compounds at various concentrations, can be used to
20 measure quantitatively the binding affinity of the laminin fragment-amyloid complex and thereby determine the effect of the test compound on the affinity between laminin fragments and amyloid. Using this information, compounds can be identified which do not modulate the binding of specific laminin fragments to amyloid and thereby allow the laminin-fragments to reduce the amyloid formation, deposition, accumulation and/or persistence,
25 and the subsequent development and persistence of amyloidosis.

Therefore a kit for practicing a method for identifying compounds useful which do not alter laminin, laminin-derived fragments or laminin-derived polypeptides to an immobilized amyloid protein, said kit comprising a) a first container having amyloid protein immobilized upon the inner surface, b) a second container which contains laminin, laminin-derived fragments or laminin-derived polypeptides dissolved in solution, c) a third container which contains antibodies specific for said laminin, laminin-derived fragments or laminin-derived polypeptides, said antibodies dissolved in solution, and d) a fourth container which contains labelled antibodies specific for laminin, laminin-derived fragments or laminin-derived polypeptides, said antibodies dissolved in solution.

10

15

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Gerardo Castillo and Alan Snow
- (ii) TITLE OF INVENTION: Therapeutic and Diagnostic Applications of Laminin and Laminin-Derived Protein Fragments
- (iii) NUMBER OF SEQUENCES: 11
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Patrick M. Dwyer
 - (B) STREET: 1919 One Union Square, 600 University Street
 - (C) CITY: Seattle
 - (D) STATE: WA (Washington)
 - (E) COUNTRY: United States of America
 - (F) ZIP: 98101
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette - 3.50 inch, 1.44 Mb storage
 - (B) COMPUTER: IBM PC
 - (C) OPERATING SYSTEM: PC-DOS (Windows NT Version 4.0, '95)
 - (D) SOFTWARE: WordPerfect 5.2
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/947,057
 - (B) FILING DATE: 08-October-1997
 - (C) CLASSIFICATION: U.S. Utility Appl.
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/027,981
 - (B) FILING DATE: 08-October-1996
 - (C) CLASSIFICATION: U.S. Provisional Appl.
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Dwyer, Patrick M.
 - (B) REGISTRATION NUMBER: 32,411
 - (C) REFERENCE/DOCKET NUMBER: PROTEO.P03
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (206) 343-7074
 - (B) TELEFAX: (206) 343-7085

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 11 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P19137

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Leu His Arg Glu His Gly Glu Leu Pro Pro Glu
1 5 10

INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 177 AMINO ACIDS
- (B) TYPE: AMINO ACID

(C) STRANDEDNESS:
(D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF
GENEBANK ACCESSION NUMBER P19137 (AMINO ACIDS #2746-2922)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Leu	Gln	Val	Gln	Leu	Ser	Ile	Arg	Thr	Phe	Ala	Ser	Ser	Gly	Leu	Ile	Tyr	Tyr	Val	Ala
1				5					10					15					20
His	Gln	Asn	Gln	Met	Asp	Tyr	Ala	Thr	Leu	Gln	Leu	Gln	Glu	Gly	Arg	Leu	His	Phe	Met
				25					30					35					40
Phe	Asp	Leu	Gly	Lys	Gly	Arg	Thr	Lys	Val	Ser	His	Pro	Ala	Leu	Leu	Ser	Asp	Gly	Lys
				45					50					55					60
Trp	His	Thr	Val	Lys	Thr	Glu	Tyr	Ile	Lys	Arg	Lys	Ala	Phe	Met	Thr	Val	Asp	Gly	Gln
				65					70					75					80
Glu	Ser	Pro	Ser	Val	Thr	Val	Val	Gly	Asn	Ala	Thr	Thr	Leu	Asp	Val	Glu	Arg	Lys	Leu
				85					90					95					100
Tyr	Leu	Gly	Gly	Leu	Pro	Ser	His	Tyr	Arg	Ala	Arg	Asn	Ile	Gly	Thr	Ile	Thr	His	Ser
				105					110					115					120
Ile	Pro	Ala	Cys	Ile	Gly	Glu	Ile	Met	Val	Asn	Gly	Gln	Gln	Leu	Asp	Lys	Asp	Arg	Pro
				125					130					135					140
Leu	Ser	Ala	Ser	Ala	Val	Asp	Arg	Cys	Tyr	Val	Val	Ala	Gln	Glu	Gly	Thr	Phe	Phe	Glu
				145					150					155					160
Gly	Ser	Gly	Tyr	Ala	Ala	Leu	Val	Lys	Glu	Gly	Tyr	Lys	Val	Arg	Leu	Asp			
				165					170					175					

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 177 AMINO ACIDS
(B) TYPE: AMINO ACID
(C) STRANDEDNESS:
(D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF
GENEBANK ACCESSION NUMBER P25391 (AMINO ACIDS #2737-2913)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Leu	Ser	Val	Glu	Leu	Ser	Ile	Arg	Thr	Phe	Ala	Ser	Ser	Gly	Leu	Ile	Tyr	Tyr	Met	Ala
1				5					10					15					20
His	Gln	Asn	Gln	Ala	Asp	Tyr	Ala	Val	Leu	Gln	Leu	His	Gly	Gly	Arg	Leu	His	Phe	Met
				25					30					35					40
Phe	Asp	Leu	Gly	Lys	Gly	Arg	Thr	Lys	Val	Ser	His	Pro	Ala	Leu	Leu	Ser	Asp	Gly	Lys
				45					50					55					60
Trp	His	Thr	Val	Lys	Thr	Asp	Tyr	Val	Lys	Arg	Lys	Gly	Phe	Ile	Thr	Val	Asp	Gly	Arg
				65					70					75					80
Glu	Ser	Pro	Met	Val	Thr	Val	Val	Gly	Asp	Gly	Thr	Met	Leu	Asp	Val	Glu	Gly	Leu	Phe
				85					90					95					100
Tyr	Leu	Gly	Gly	Leu	Pro	Ser	Gln	Tyr	Gln	Ala	Arg	Lys	Ile	Gly	Asn	Ile	Thr	His	Ser
				105					110					115					120
Ile	Pro	Ala	Cys	Ile	Gly	Asp	Val	Thr	Val	Asn	Ser	Lys	Gln	Leu	Asp	Lys	Asp	Ser	Pro
				125					130					135					140
Val	Ser	Ala	Phe	Thr	Val	Asn	Arg	Cys	Tyr	Ala	Val	Ala	Gln	Glu	Gly	Thr	Tyr	Phe	Asp

	145	150	155	160
Gly Ser Gly Tyr	Ala Ala Leu Val Lys	Glu Gly Tyr Lys Val	Gln Ser Asp	
	165	170	175	

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3084 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P19137

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met	Arg	Gly	Ser	Gly	Thr	Gly	Ala	Ala	Leu	Leu	Val	Leu	Leu	Ala	Ser	Val	Leu	Trp	Val
1				5					10					15					20
Thr	Val	Arg	Ser	Gln	Gln	Arg	Gly	Leu	Phe	Pro	Ala	Ile	Leu	Asn	Leu	Ala	Thr	Asn	Ala
				25					30					35					40
His	Ile	Ser	Ala	Asn	Ala	Thr	Cys	Gly	Glu	Lys	Gly	Pro	Glu	Met	Phe	Cys	Lys	Leu	Val
				45					50					55					60
Glu	His	Val	Pro	Gly	Arg	Pro	Val	Arg	His	Ala	Gln	Cys	Arg	Val	Cys	Asp	Gly	Asn	Ser
				65					70					75					80
Thr	Asn	Pro	Arg	Glu	Arg	His	Pro	Ile	Ser	His	Ala	Ile	Asp	Gly	Thr	Asn	Asn	Trp	Trp
				85					90					95					100
Gln	Ser	Pro	Ser	Ile	Gln	Asn	Gly	Arg	Glu	Tyr	His	Trp	Val	Thr	Val	Thr	Leu	Asp	Leu
				105					110					115					120
Arg	Gln	Val	Phe	Gln	Val	Ala	Tyr	Ile	Ile	Ile	Lys	Ala	Ala	Asn	Ala	Pro	Arg	Pro	Gly
				125					130					135					140
Asn	Trp	Ile	Leu	Glu	Arg	Ser	Val	Asp	Gly	Val	Lys	Phe	Lys	Pro	Trp	Gln	Tyr	Tyr	Ala
				145					150					155					160
Val	Ser	Asp	Thr	Glu	Cys	Leu	Thr	Arg	Tyr	Lys	Ile	Thr	Pro	Arg	Arg	Gly	Pro	Pro	Thr
				165					170					175					180
Tyr	Arg	Ala	Asp	Asn	Glu	Val	Ile	Cys	Thr	Ser	Tyr	Tyr	Ser	Lys	Leu	Val	Pro	Leu	Glu
				185					190					195					200
His	Gly	Glu	Ile	His	Thr	Ser	Leu	Ile	Asn	Gly	Arg	Pro	Ser	Ala	Asp	Asp	Pro	Ser	Pro
				205					210					215					220
Gln	Leu	Leu	Glu	Phe	Thr	Ser	Ala	Arg	Tyr	Ile	Arg	Leu	Arg	Leu	Gln	Arg	Ile	Arg	Thr
				225					230					235					240
Leu	Asn	Ala	Asp	Leu	Met	Thr	Leu	Ser	His	Arg	Asp	Leu	Arg	Asp	Leu	Asp	Pro	Ile	Val
				245					250					255					260
Thr	Arg	Arg	Tyr	Tyr	Tyr	Ser	Ile	Lys	Asp	Ile	Ser	Val	Gly	Gly	Met	Cys	Ile	Cys	Tyr
				265					270					275					280
Gly	His	Ala	Ser	Ser	Cys	Pro	Trp	Asp	Glu	Glu	Ala	Lys	Gln	Leu	Gln	Cys	Gln	Cys	Glu
				285					290					295					300
His	Asn	Thr	Cys	Gly	Glu	Ser	Cys	Asp	Arg	Cys	Cys	Pro	Gly	Tyr	His	Gln	Gln	Pro	Trp
				305					310					315					320
Arg	Pro	Gly	Thr	Ile	Ser	Ser	Gly	Asn	Glu	Cys	Glu	Glu	Cys	Asn	Cys	His	Asn	Lys	Ala
				325					330					335					340
Lys	Asp	Cys	Tyr	Tyr	Asp	Ser	Ser	Val	Ala	Lys	Glu	Arg	Arg	Ser	Leu	Asn	Thr	Ala	Gly
				345					350					355					360
Gln	Tyr	Ser	Gly	Gly	Gly	Val	Cys	Val	Asn	Cys	Ser	Gln	Asn	Thr	Thr	Gly	Ile	Asn	Cys
				365					370					375					380
Glu	Thr	Cys	Ile	Asp	Gln	Tyr	Tyr	Arg	Pro	His	Lys	Val	Ser	Pro	Tyr	Asp	Asp	His	Pro
				385					390					395					400
Cys	Arg	Pro	Cys	Asn	Cys	Asp	Pro	Val	Gly	Ser	Leu	Ser	Ser	Val	Cys	Ile	Lys	Asp	Asp

Gly	Leu	Asp	Pro	Glu	Gln	Gly	Cys	Gln	Ala	Cys	Asn	Cys	Ser	Ala	Val	Gly	Ser	Thr	Ser
				1045					1050					1055					1060
Ala	Gln	Cys	Asp	Val	Leu	Ser	Gly	His	Cys	Pro	Cys	Lys	Lys	Gly	Phe	Gly	Gly	Gln	Ser
				1065					1070					1075					1080
Cys	His	Gln	Cys	Ser	Leu	Gly	Tyr	Arg	Ser	Phe	Pro	Asp	Cys	Val	Pro	Cys	Gly	Cys	Asp
				1085					1090					1095					1100
Leu	Arg	Gly	Thr	Leu	Pro	Asp	Thr	Cys	Asp	Leu	Glu	Gln	Gly	Leu	Cys	Ser	Cys	Ser	Glu
				1105					1110					1115					1120
Asp	Ser	Gly	Thr	Cys	Ser	Cys	Lys	Glu	Asn	Val	Val	Gly	Pro	Gln	Cys	Ser	Lys	Cys	Gln
				1125					1130					1135					1140
Ala	Gly	Thr	Phe	Ala	Leu	Arg	Gly	Asp	Asn	Pro	Gln	Gly	Cys	Ser	Pro	Cys	Phe	Cys	Phe
				1145					1150					1155					1160
Gly	Leu	Ser	Gln	Leu	Cys	Ser	Glu	Leu	Glu	Gly	Tyr	Val	Arg	Thr	Leu	Ile	Thr	Leu	Ala
				1165					1170					1175					1180
Ser	Asp	Gln	Pro	Leu	Leu	His	Val	Val	Ser	Gln	Ser	Asn	Leu	Lys	Gly	Thr	Ile	Glu	Gly
				1185					1190					1195					1200
Val	His	Phe	Gln	Pro	Pro	Asp	Thr	Leu	Leu	Asp	Ala	Glu	Ala	Val	Arg	Gln	His	Ile	Tyr
				1205					1210					1215					1220
Ala	Glu	Pro	Phe	Tyr	Trp	Arg	Leu	Pro	Lys	Gln	Phe	Gln	Gly	Asp	Gln	Leu	Leu	Ala	Tyr
				1225					1230					1235					1240
Gly	Gly	Lys	Leu	Gln	Tyr	Ser	Val	Ala	Phe	Tyr	Ser	Thr	Leu	Gly	Thr	Gly	Thr	Ser	Asn
				1245					1250					1255					1260
Tyr	Glu	Pro	Gln	Val	Leu	Ile	Lys	Gly	Gly	Arg	Ala	Arg	Lys	His	Val	Ile	Tyr	Met	Asp
				1265					1270					1275					1280
Ala	Pro	Ala	Pro	Glu	Asn	Gly	Val	Arg	Gln	Asp	Tyr	Glu	Val	Gln	Met	Lys	Glu	Glu	Phe
				1285					1290					1295					1300
Trp	Lys	Tyr	Phe	Asn	Ser	Val	Ser	Glu	Lys	His	Val	Thr	His	Ser	Asp	Phe	Met	Ser	Val
				1305					1310					1315					1320
Leu	Ser	Asn	Ile	Asp	Tyr	Ile	Leu	Ile	Lys	Ala	Ser	Tyr	Gly	Gln	Gly	Leu	Gln	Gln	Ser
				1325					1330					1335					1340
Arg	Ile	Ala	Asn	Ile	Ser	Met	Glu	Val	Gly	Arg	Lys	Ala	Val	Glu	Leu	Pro	Ala	Glu	Gly
				1345					1350					1355					1360
Glu	Ala	Ala	Leu	Leu	Leu	Glu	Leu	Cys	Val	Cys	Pro	Pro	Gly	Thr	Ala	Gly	His	Ser	Cys
				1365					1370					1375					1380
Gln	Asp	Cys	Ala	Pro	Gly	Tyr	Tyr	Arg	Glu	Lys	Leu	Pro	Glu	Ser	Gly	Gly	Arg	Gly	Pro

Lys Val Ala Thr	1665	1670	1675	1680
Gln Ser Met His	1685	1690	1695	1700
Glu Met Gln Gln	1705	1710	1715	1720
Gln Lys Arg Phe	1725	1730	1735	1740
Leu Ser Asn His	1745	1750	1755	1760
Lys Thr Gln Glu	1765	1770	1775	1780
Gln Glu Lys Lys	1785	1790	1795	1800
Gly Arg Glu Trp	1805	1810	1815	1820
Gln Leu Glu His	1825	1830	1835	1840
Asp Asp Leu Val	1845	1850	1855	1860
Gln His Ala Ser	1865	1870	1875	1880
Arg Asn Val Ser	1885	1890	1895	1900
Thr Glu Glu Ala	1905	1910	1915	1920
Ile Ser Glu Ser	1925	1930	1935	1940
Lys Glu Ser Val	1945	1950	1955	1960
Lys Asn Leu Thr	1965	1970	1975	1980
Ser Leu Ala Met	1985	1990	1995	2000
Glu Leu Ala Ala	2005	2010	2015	2020
Ser Leu Arg Val	2025	2030	2035	2040
Thr Asn Asp Leu	2045	2050	2055	2060
Asp Met Glu Met	2065	2070	2075	2080
Glu Asn Leu Ser	2085	2090	2095	2100
Ala Ala Ser Ile	2105	2110	2115	2120
Gln Thr Ser Ser	2125	2130	2135	2140
Asn Leu Leu Phe	2145	2150	2155	2160
Arg Gly Lys Val	2165	2170	2175	2180
Glu Val Ser Ile	2185	2190	2195	2200
Gly Ser Leu Ser	2205	2210	2215	2220
Ser Pro Gly Pro	2225	2230	2235	2240
Leu Gly Gly Gln	2245	2250	2255	2260
Gly Glu Ala Phe	2265	2270	2275	2280
	2285	2290	2295	2300

Lys Cys Asn Gly	Cys Phe Gly Ser Ser	Gln Asn Glu Asp Ser	Ser Phe His Phe Asp Gly
2305	2310	2315	2320
Ser Gly Tyr Ala	Met Val Glu Lys Thr	Leu Arg Pro Thr Val	Thr Gln Ile Val Ile Leu
2325	2330	2335	2340
Phe Ser Thr Phe	Ser Pro Asn Gly Leu	Leu Phe Tyr Leu Ala	Ser Asn Gly Thr Lys Asp
2345	2350	2355	2360
Phe Leu Ser Ile	Glu Leu Val Arg Gly	Arg Val Lys Val Met	Val Asp Leu Gly Ser Gly
2365	2370	2375	2380
Pro Leu Thr Leu	Met Thr Asp Arg Arg	Tyr Asn Asn Gly Thr	Trp Tyr Lys Ile Ala Phe
2385	2390	2395	2400
Gln Arg Asn Arg	Lys Gln Gly Leu Leu	Ala Val Phe Asp Ala	Tyr Asp Thr Ser Asp Lys
2405	2410	2415	2420
Glu Thr Lys Gln	Gly Glu Thr Pro Gly	Ala Ala Ser Asp Leu	Asn Arg Leu Glu Lys Asp
2425	2430	2435	2440
Leu Ile Tyr Val	Gly Gly Leu Pro His	Ser Lys Ala Val Arg	Lys Gly Val Ser Ser Arg
2445	2450	2455	2460
Ser Tyr Val Gly	Cys Ile Lys Asn Leu	Glu Ile Ser Arg Ser	Thr Phe Asp Leu Leu Arg
2465	2470	2475	2480
Asn Ser Tyr Gly	Val Arg Lys Gly Cys	Ala Leu Glu Pro Ile	Gln Ser Val Ser Phe Leu
2485	2490	2495	2500
Arg Gly Gly Tyr	Val Glu Met Pro Pro	Lys Ser Leu Ser Pro	Glu Ser Ser Leu Leu Ala
2505	2510	2515	2520
Thr Phe Ala Thr	Lys Asn Ser Ser Gly	Ile Leu Leu Val Ala	Leu Gly Lys Asp Ala Glu
2525	2530	2535	2540
Glu Ala Gly Gly	Ala Gln Ala His Val	Pro Phe Phe Ser Ile	Met Leu Leu Glu Gly Arg
2545	2550	2555	2560
Ile Glu Val His	Val Asn Ser Gly Asp	Gly Thr Ser Leu Arg	Lys Ala Leu Leu His Ala
2565	2570	2575	2580
Pro Thr Gly Ser	Tyr Ser Asp Gly Gln	Glu His Ser Ile Ser	Leu Val Arg Asn Arg Arg
2585	2590	2595	2600
Val Ile Thr Ile	Gln Val Asp Glu Asn	Ser Pro Val Glu Met	Lys Leu Gly Pro Leu Thr
2605	2610	2615	2620
Glu Gly Lys Thr	Ile Asp Ile Ser Asn	Leu Tyr Ile Gly Gly	Leu Pro Glu Asp Lys Ala
2625	2630	2635	2640
Thr Pro Met Leu	Lys Met Arg Thr Ser	Phe His Gly Cys Ile	Lys Asn Val Val Leu Asp
2645	2650	2655	2660
Ala Gln Leu Leu	Asp Phe Thr His Ala	Thr Gly Ser Glu Gln	Val Glu Leu Asp Thr Cys
2665	2670	2675	2680
Leu Leu Ala Glu	Glu Pro Met Gln Ser	Leu His Arg Glu His	Gly Glu Leu Pro Pro Glu
2685	2690	2695	2700
Pro Pro Thr Leu	Pro Gln Pro Glu Leu	Cys Ala Val Asp Thr	Ala Pro Gly Tyr Val Ala
2705	2710	2715	2720
Gly Ala His Gln	Phe Gly Leu Ser Gln	Asn Ser His Leu Val	Leu Pro Leu Asn Gln Ser
2725	2730	2735	2740
Asp Val Arg Lys	Arg Leu Gln Val Gln	Leu Ser Ile Arg Thr	Phe Ala Ser Ser Gly Leu
2745	2750	2755	2760
Ile Tyr Tyr Val	Ala His Gln Asn Gln	Met Asp Tyr Ala Thr	Leu Gln Leu Gln Glu Gly
2765	2770	2775	2780
Arg Leu His Phe	Met Phe Asp Leu Gly	Lys Gly Arg Thr Lys	Val Ser His Pro Ala Leu
2785	2790	2795	2800
Leu Ser Asp Gly	Lys Trp His Thr Val	Lys Thr Glu Tyr Ile	Lys Arg Lys Ala Phe Met
2805	2810	2815	2820
Thr Val Asp Gly	Gln Glu Ser Pro Ser	Val Thr Val Val Gly	Asn Ala Thr Thr Leu Asp
2825	2830	2835	2840
Val Glu Arg Lys	Leu Tyr Leu Gly Gly	Leu Pro Ser His Tyr	Arg Ala Arg Asn Ile Gly
2845	2850	2855	2860
Thr Ile Thr His	Ser Ile Pro Ala Cys	Ile Gly Glu Ile Met	Val Asn Gly Gln Gln Leu
2865	2870	2875	2880
Asp Lys Asp Arg	Pro Leu Ser Ala Ser	Ala Val Asp Arg Cys	Tyr Val Val Ala Gln Glu
2885	2890	2895	2900
Gly Thr Phe Phe	Glu Gly Ser Gly Tyr	Ala Ala Leu Val Lys	Glu Gly Tyr Lys Val Arg
2905	2910	2915	2920
Leu Asp Leu Asn	Ile Thr Leu Glu Phe	Arg Thr Thr Ser Lys	Asn Gly Val Leu Leu Gly

Ile Ser Ser Ala	2925	Lys Val Asp Ala Ile	2930	Gly Leu Glu Ile Val	2935	Asp Gly Lys Val Leu	2940
	2945		2950		2955		2960
His Val Asn Asn	2965	Gly Ala Gly Arg Ile	2970	Thr Ala Thr Tyr Gln	2975	Pro Arg Ala Ala Arg	2980
Leu Cys Asp Gly	2985	Lys Trp His Thr Leu	2990	Gln Ala His Lys Ser	2995	Lys His Arg Ile Val	3000
Thr Val Asp Gly	3005	Asn Ser Val Arg Ala	3010	Glu Ser Pro His Thr	3015	His Ser Thr Ser Ala	3020
Thr Asn Asp Pro	3025	Ile Tyr Val Gly Gly	3030	Tyr Pro Ala His Ile	3035	Lys Gln Asn Cys Leu	3040
Ser Arg Ala Ser	3045	Phe Arg Gly Cys Val	3050	Arg Asn Leu Arg Leu	3055	Ser Arg Gly Ser Gln	3060
Gln Ser Leu Asp	3065	Leu Ser Arg Ala Phe	3070	Asp Leu Gln Gly Val	3075	Phe Pro His Ser Cys	3080
Gly Pro Glu Pro							

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3075 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF
GENEBANK ACCESSION NUMBER P25391

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Met Arg Gly Gly Val Leu Leu Val Leu	1	Leu Leu Cys Val Ala Ala Gln Cys Arg Gln Arg	20
Gly Leu Phe Pro Ala Ile Leu Asn Leu	5	Ala Ser Asn Ala His Ile Ser Thr Asn Ala Thr	35
Cys Gly Glu Lys Gly Pro Glu Met Phe	10	Cys Lys Leu Val Glu His Val Pro Gly Arg Pro	40
Val Arg Asn Pro Gln Cys Arg Ile Cys	15	Asn Pro Arg Glu Arg His	55
Pro Ile Ser His Ala Ile Asp Gly Thr	20	Asp Gly Asn Ser Ala Asn Pro Arg Glu Arg His	60
Gly Arg Glu Tyr His Trp Val Thr Ile	25	Asn Asn Trp Trp Gln Ser Pro Ser Ile Gln Asn	75
Tyr Val Ile Ile Lys Ala Ala Asn Ala	30	Thr Leu Asp Leu Arg Gln Val Phe Gln Val Ala	90
Leu Asp Gly Thr Thr Phe Ser Pro Trp	35	Pro Arg Pro Gly Asn Trp Ile Leu Glu Arg Ser	105
Ser Arg Tyr Asn Ile Thr Pro Arg Arg	40	Gln Tyr Tyr Ala Val Ser Asp Ser Glu Cys Leu	120
Ile Cys Thr Ser Tyr Tyr Ser Arg Leu	45	Val Pro Leu Glu His Gly Glu Ile His Thr Ser	135
Leu Ile Asn Gly Arg Pro Ser Ala Asp	50	Arg Gly Pro Pro Thr Tyr Arg Ala Asp Asp Glu Val	150
Ala Arg Tyr Ile Arg Leu Arg Leu Gln	55	Val Pro Leu Glu His Gly Glu Ile His Thr Ser	165
Leu Ser His Arg Glu Pro Lys Glu Leu	60	Asp Leu Ser Pro Lys Leu Leu Glu Phe Thr Ser	180
Ile Lys Asp Ile Ser Val Gly Gly Met	65	Asp Leu Ser Pro Lys Leu Leu Glu Phe Thr Ser	195
Trp Asp Glu Thr Thr Lys Lys Leu Gln	70	Cys Ile Cys Tyr Gly His Ala Ser Ser Cys Pro	210
	75	Cys Gln Cys Glu His Asn Thr Cys Gly Glu Ser	225
	80		240
	85		255
	90		270
	95		285
	100		300
	105		315
	110		330
	115		345
	120		360
	125		375
	130		390
	135		405
	140		420
	145		435
	150		450
	155		465
	160		480
	165		495
	170		510
	175		525
	180		540
	185		555
	190		570
	195		585
	200		600
	205		615
	210		630
	215		645
	220		660
	225		675
	230		690
	235		705
	240		720
	245		735
	250		750
	255		765
	260		780
	265		795
	270		810
	275		825
	280		840
	285		855
	290		870
	295		885
	300		900
	305		915
	310		930
	315		945
	320		960
	325		975
	330		990
	335		1005
	340		1020
	345		1035
	350		1050
	355		1065
	360		1080
	365		1095
	370		1110
	375		1125
	380		1140
	385		1155
	390		1170
	395		1185
	400		1200
	405		1215
	410		1230
	415		1245
	420		1260
	425		1275
	430		1290
	435		1305
	440		1320
	445		1335
	450		1350
	455		1365
	460		1380
	465		1395
	470		1410
	475		1425
	480		1440
	485		1455
	490		1470
	495		1485
	500		1500
	505		1515
	510		1530
	515		1545
	520		1560
	525		1575
	530		1590
	535		1605
	540		1620
	545		1635
	550		1650
	555		1665
	560		1680
	565		1695
	570		1710
	575		1725
	580		1740
	585		1755
	590		1770
	595		1785
	600		1800
	605		1815
	610		1830
	615		1845
	620		1860
	625		1875
	630		1890
	635		1905
	640		1920
	645		1935
	650		1950
	655		1965
	660		1980
	665		1995
	670		2010
	675		2025
	680		2040
	685		2055
	690		2070
	695		2085
	700		2100
	705		2115
	710		2130
	715		2145
	720		2160
	725		2175
	730		2190
	735		2205
	740		2220
	745		2235
	750		2250
	755		2265
	760		2280
	765		2295
	770		2310
	775		2325
	780		2340
	785		2355
	790		2370
	795		2385
	800		2400
	805		2415
	810		2430
	815		2445
	820		2460
	825		2475
	830		2490
	835		2505
	840		2520
	845		2535
	850		2550
	855		2565
	860		2580
	865		2595
	870		2610
	875		2625
	880		2640
	885		2655
	890		2670
	895		2685
	900		2700
	905		2715
	910		2730
	915		2745
	920		2760
	925		2775
	930		2790
	935		2805
	940		2820
	945		2835
	950		2850
	955		2865
	960		2880
	965		2895
	970		2910
	975		2925
	980		2940
	985		2955
	990		2970
	995		2985
	1000		2995
	1005		3005
	1010		3015
	1015		3025
	1020		3035
	1025		3045
	1030		3055
	1035		3065
	1040		3075
	1045		
	1050		
	1055		
	1060		
	1065		
	1070		
	1075		
	1080		
	1085		
	1090		
	1095		
	1100		
	1105		
	1110		
	1115		
	1120		
	1125		
	1130		
	1135		
	1140		
	1145		
	1150		
	1155		
	1160		
	1165		
	1170		
	1175		
	1180		
	1185		
	1190		
	1195		
	1200		
	1205		
	1210		
	1215		
	1220		
	1225		
	1230		
	1235		
	1240		
	1245		
	1250		
	1255		
	1260		
	1265		
	1270		
	1275		
	1280		
	1285		
	1290		
	1295		
	1300		
	1305		
	1310		
	1315		
	1320		
	1325		
	1330		
	1335		
	1340		
	1345		
	1350		
	1355		
	1360		
	1365		
	1370		
	1375		
	1380		
	1385		
	1390		
	1395		
	1400		
	1405		
	1410		
	1415		
	1420		
	1425		
	1430		
	1435		
	1440		
	1445		
	1450		
	1455		
	1460		
	1465		
	1470		
	1475		
	1480		
	1485		
	1490		
	1495		
	1500		
	1505		
	1510		
	1515		
	1520		
	1525		
	1530		
	1535		
	1540		
	1545		

Cys Asn Arg Cys	285	Cys Pro Gly Tyr His	290	Pro Gly Thr Val Ser Ser	295	300
Gly Asn Thr Cys	305	Glu Ala Cys Asn Cys	310	Asp Cys Tyr Tyr Asp Glu	315	320
Ser Val Ala Lys	325	Gln Lys Lys Ser Leu	330	Phe Arg Gly Gly Gly Tyr	335	340
Cys Ile Asn Cys	345	Leu Gln Asn Thr Met	350	Thr Cys Ile Asp Gly Tyr	355	360
Tyr Arg Pro His	365	Lys Val Ser Pro Tyr	370	Arg Pro Cys Asn Cys Asp	375	380
Pro Val Gly Ser	385	Leu Ser Ser Val Cys	390	His Ser Asp Leu His Asn	395	400
Gly Lys Gln Pro	405	Gly Gln Cys Pro Cys	410	Lys Asp Asp Leu His Ser	415	420
Cys Gln Leu Gly	425	Tyr Lys Asp Tyr Pro	430	Gly Glu Lys Cys Asp Arg	435	440
Ser Ala Ser Asp	445	Glu Pro Cys Thr Gly	450	Gly Cys Asn Pro Val Gly	455	460
Ala Cys Asp Arg	465	Cys Lys Pro Gly Phe	470	Glu Asn Val Glu Gly Lys	475	480
Ser Glu Cys Phe	485	Cys Phe Gly Val Ser	490	Lys Asn Pro Arg Gly Cys	495	500
Gln Val Asn Ser	505	Met Ser Gly Trp Leu	510	Leu Ser Trp Pro Val Gly	515	520
Ser Gln Gln Asp	525	Ala Leu Gly Gly Arg	530	Ser Pro Arg Lys Ile Pro	535	540
Gln Arg Leu Ala	545	Pro Lys Tyr Tyr Trp	550	Asn Asn Thr Ala Val Met	555	560
Thr Ala Phe Gly	565	Gly Phe Leu Lys Tyr	570	Tyr Leu Gly Asn Lys Leu	575	580
Asp Ser Asn Leu	585	Met Ser His Ala Asp	590	Ile Pro Val Glu Thr Val	595	600
Thr Gln Ala Glu	605	Gly Leu Ser Leu Gln	610	Asn Gly Leu Thr Leu Ser	615	620
Val Pro Glu Asn	625	Phe Gln Asp Phe His	630	Leu Asn Val Val Arg Leu	635	640
Thr Val Leu Ala	645	Asn Val Thr His Leu	650	Asp Arg Asp Gln Leu Met	655	660
Ala Leu Tyr Arg	665	Leu Glu Ser Val Ser	670	Tyr Asn Ser Ala Lys Met	675	680
Val Val Ala Ala	685	Asp Val Glu His Cys	690	Ser Asn Ala Ile Asp Leu	695	700
Glu Ser Cys Leu	705	Ser Gly Tyr Tyr Arg	710	Gly Thr Gly Thr Ser Cys	715	720
Pro Cys Glu Cys	725	His Gly His Ala Ala	730	Phe Gly Gly Ile Cys Gln	735	740
Ala His Asn Thr	745	Thr Gly Val His Cys	750	Gly Val Cys Ile Ala Cys	755	760
Ser Arg Gly Thr	765	Pro Gly Asp Cys Gln	770	Gly Phe Tyr Gly Glu Pro	775	780
Asn Phe Ser Pro	785	Thr Cys His Leu Asn	790	Leu Thr Ile Ala Ser Asn	795	800
Pro Gly Tyr Ser	805	Gly Ala Trp Cys Glu	810	Val Cys Asp Trp Cys Ala	815	820
Val Pro Gly Glu	825	Ser Cys Val Pro Cys	830	Tyr Tyr Gly Asn Pro Thr	835	840
Gly His Cys Asp	845	Ser Val Thr Gly Glu	850	Val Asp Pro Ser Glu Ala	855	860
His Cys Glu Arg	865	Cys Ala Asp Gly Phe	870	Gly Asn Thr Asp Gly Ala	875	880
Ala Cys Glu Cys	885	His Val Lys Gly Ser	890	Thr Ala Lys Asn Cys Arg	895	900
	905		910	His Leu Glu Thr Gly Leu	915	920

Cys Asp Cys Lys	Pro Asn Val Thr Gly	Gln Gln Cys Asp Gln	Cys Leu His Gly Tyr Tyr
	925	930	935
Gly Leu Asp Ser	Gly His Gly Cys Arg	Pro Cys Asn Cys Ser	Val Ala Gly Ser Val Ser
	945	950	955
Asp Gly Cys Thr	Asp Glu Gly Gln Cys	His Cys Val Pro Gly	Val Ala Gly Lys Arg Cys
	965	970	975
Asp Arg Cys Ala	His Gly Phe Tyr Ala	Tyr Gln Asp Gly Ser	Cys Thr Pro Cys Asp Cys
	985	990	995
Pro His Thr Gln	Asn Thr Cys Asp Pro	Glu Thr Gly Glu Cys	Val Cys Pro Pro His Thr
	1005	1010	1015
Gln Gly Gly Lys	Cys Glu Glu Cys Glu	Asp Gly His Trp Gly	Tyr Asp Ala Glu Val Gly
	1025	1030	1035
Cys Gln Ala Cys	Asn Cys Ser Leu Val	Gly Ser Thr His His	Arg Cys Asp Val Val Thr
	1045	1050	1055
Gly His Cys Gln	Cys Lys Ser Lys Phe	Gly Gly Arg Ala Cys	Asp Gln Cys Ser Leu Gly
	1065	1070	1075
Tyr Arg Asp Phe	Pro Asp Cys Val Pro	Cys Asp Cys Asp Leu	Arg Gly Thr Ser Gly Asp
	1085	1090	1095
Ala Cys Asn Leu	Glu Gln Gly Leu Cys	Gly Cys Val Glu Glu	Thr Gly Ala Cys Pro Cys
	1105	1110	1115
Lys Glu Asn Val	Phe Gly Pro Gln Cys	Asn Glu Cys Arg Glu	Gly Thr Phe Ala Leu Arg
	1125	1130	1135
Ala Asp Asn Pro	Leu Gly Cys Ser Pro	Cys Phe Cys Ser Gly	Leu Ser His Leu Cys Ser
	1145	1150	1155
Glu Leu Glu Asp	Tyr Val Arg Thr Pro	Val Thr Leu Gly Ser	Asp Gln Pro Leu Leu Arg
	1165	1170	1175
Val Val Ser Gln	Ser Asn Leu Arg Gly	Thr Thr Glu Gly Val	Tyr Tyr Gln Ala Pro Asp
	1185	1190	1195
Phe Leu Leu Asp	Ala Ala Thr Val Arg	Gln His Ile Arg Ala	Glu Pro Phe Tyr Trp Arg
	1205	1210	1215
Leu Pro Gln Gln	Phe Gln Gly Asp Gln	Leu Met Ala Tyr Gly	Gly Lys Leu Lys Tyr Ser
	1225	1230	1235
Val Ala Phe Tyr	Ser Leu Asp Gly Val	Gly Thr Ser Asn Phe	Glu Pro Gln Val Leu Ile
	1245	1250	1255
Lys Gly Gly Arg	Ile Arg Lys Gln Val	Ile Tyr Met Asp Ala	Pro Ala Pro Glu Asn Gly
	1265	1270	1275
Val Arg Gln Glu	Gln Glu Val Ala Met	Arg Glu Asn Phe Trp	Lys Tyr Phe Asn Ser Val
	1285	1290	1295
Ser Glu Lys Pro	Val Thr Arg Glu Asp	Phe Met Ser Val Leu	Ser Asp Ile Glu Tyr Ile
	1305	1310	1315
Leu Ile Lys Ala	Ser Tyr Gly Gln Gly	Leu Gln Gln Ser Arg	Ile Ser Asp Ile Ser Val
	1325	1330	1335
Glu Val Gly Arg	Lys Ala Glu Lys Leu	His Pro Glu Glu Glu	Val Ala Ser Leu Leu Glu
	1345	1350	1355
Asn Cys Val Cys	Pro Pro Gly Thr Val	Gly Phe Ser Cys Gln	Asp Cys Ala Pro Gly Tyr
	1365	1370	1375
His Arg Gly Lys	Leu Pro Ala Gly Ser	Asp Arg Gly Pro Arg	Pro Leu Val Ala Pro Cys
	1385	1390	1395
Val Pro Cys Ser	Cys Asn Asn His Ser	Asp Thr Cys Asp Pro	Asn Thr Gly Lys Cys Leu
	1405	1410	1415
Asn Cys Gly Asp	Asn Thr Ala Gly Asp	His Cys Asp Val Cys	Thr Ser Gly Tyr Tyr Gly
	1425	1430	1435
Lys Val Thr Gly	Ser Ala Ser Asp Cys	Ala Leu Cys Ala Cys	Pro His Ser Pro Pro Ala
	1445	1450	1455
Ser Phe Ser Pro	Thr Cys Val Leu Glu	Gly Asp His Asp Phe	Arg Cys Asp Ala Cys Leu
	1465	1470	1475
Leu Gly Tyr Glu	Gly Lys His Cys Glu	Arg Cys Ser Ser Ser	Tyr Tyr Gly Asn Pro Gln
	1485	1490	1495
Thr Pro Gly Gly	Ser Cys Gln Lys Cys	Asp Cys Asn Arg His	Gly Ser Val His Gly Asp
	1505	1510	1515
Cys Asp Arg Thr	Ser Gly Gln Cys Val	Cys Arg Leu Gly Ala	Ser Gly Leu Arg Cys Asp
	1525	1530	1535
Glu Cys Glu Pro	Arg His Ile Leu Met	Glu Thr Asp Cys Val	Ser Cys Asp Asp Glu Cys

Val Gly Val Leu	1545	Leu Asn Asp Leu Asp	1550	Glu Ile Gly Asp Ala	1555	Val Leu Ser Leu Asn	1560
Thr Gly Ile Ile	1565	Pro Val Pro Tyr Gly	1570	Ile Leu Ser Asn Leu	1575	Glu Asn Thr Thr Lys	1580
Leu Gln Glu Ser	1585	Leu Leu Lys Glu Asn	1590	Met Gln Lys Asp Leu	1595	Gly Lys Ile Lys Leu	1600
Gly Val Ala Glu	1605	Glu Thr Asp Asn Leu	1610	Gln Lys Lys Leu Thr	1615	Arg Met Leu Ala Ser	1620
Gln Lys Val Asn	1625	Arg Ala Thr Glu Arg	1630	Ile Phe Lys Glu Ser	1635	Gln Asp Leu Ala Val	1640
Ile Glu Arg Leu	1645	Gln Met Ser Ile Thr	1650	Glu Ile Met Glu Lys	1655	Thr Thr Leu Asn Gln	1660
Leu Asp Glu Asp	1665	Phe Leu Leu Pro Asn	1670	Ser Thr Leu Gln Asn	1675	Met Gln Gln Asn Gly	1680
Ser Leu Leu Glu	1685	Ile Met Gln Ile Arg	1690	Asp Phe Thr Gln Leu	1695	His Gln Asn Ala Thr	1700
Glu Leu Lys Ala	1705	Ala Glu Asp Leu Leu	1710	Ser Gln Ile Gln Glu	1715	Asn Tyr Gln Lys Pro	1720
Glu Glu Leu Glu	1725	Val Leu Lys Glu Ala	1730	Ala Ser His Val Leu	1735	Ser Lys His Asn Asn	1740
Leu Lys Ala Ala	1745	Glu Ala Leu Val Arg	1750	Glu Ala Glu Ala Lys	1755	Met Gln Glu Ser Asn	1760
Leu Leu Leu Met	1765	Val Asn Ala Asn Leu	1770	Arg Glu Phe Ser Asp	1775	Lys Lys Leu His Val	1780
Glu Glu Gln Asn	1785	Leu Thr Ser Glu Leu	1790	Ile Val Gln Gly Arg	1795	Gly Leu Ile Asp Ala	1800
Ala Ala Gln Thr	1805	Asp Ala Val Gln Asp	1810	Ala Leu Glu His Leu	1815	Glu Asp His Gln Asp	1820
Leu Leu Leu Trp	1825	Ser Ala Lys Ile Arg	1830	His His Ile Asp Asp	1835	Leu Val Met His Met	1840
Gln Arg Asn Ala	1845	Val Asp Leu Val Tyr	1850	Arg Ala Glu Asp His	1855	Ala Thr Glu Phe Gln	1860
Leu Ala Asp Val	1865	Leu Tyr Ser Gly Leu	1870	Glu Asn Ile Arg Asn	1875	Val Ser Leu Asn Ala	1880
Ser Ala Ala Tyr	1885	Val His Tyr Asn Ile	1890	Gln Ser Leu Ile Glu	1895	Glu Ser Glu Glu Leu	1900
Arg Asp Ala His	1905	Arg Thr Val Thr Glu	1910	Thr Ser Leu Leu Ser	1915	Glu Ser Leu Val Ser	1920
Gly Lys Ala Ala	1925	Val Gln Arg Ser Ser	1930	Arg Phe Leu Lys Glu	1935	Gly Asn Asn Leu Ser	1940
Lys Leu Pro Gly	1945	Ile Ala Leu Glu Leu	1950	Ser Glu Leu Arg Asn	1955	Lys Thr Asn Arg Phe	1960
Glu Asn Ala Val	1965	Glu Ile Thr Arg Gln	1970	Thr Asn Glu Ser Leu	1975	Leu Ile Leu Arg Ala	1980
Pro Glu Gly Ile	1985	Arg Asp Lys Gly Ala	1990	Lys Thr Lys Glu Leu	1995	Ala Thr Ser Ala Ser	2000
Ser Ala Val Ser	2005	Thr Leu Arg Asp Val	2010	Ala Gly Leu Ser Gln	2015	Glu Leu Leu Asn Thr	2020
Ala Ser Leu Ser	2025	Arg Val Asn Thr Thr	2030	Leu Arg Glu Thr His	2035	Gln Leu Leu Gln Asp	2040
Thr Met Ala Thr	2045	Leu Leu Ala Gly Arg	2050	Lys Val Lys Asp Val	2055	Glu Ile Gln Ala Asn	2060
Leu Phe Asp Arg	2065	Leu Lys Pro Leu Lys	2070	Met Leu Glu Glu Asn	2075	Leu Ser Arg Asn Leu	2080
Glu Ile Lys Leu	2085	Leu Ile Ser Gln Ala	2090	Arg Lys Gln Ala Ala	2095	Ser Ile Lys Val Ala	2100
Ser Ala Asp Arg	2105	Asp Cys Ile Arg Ala	2110	Tyr Gln Pro Gln Ile	2115	Ser Ser Thr Asn Tyr	2120
Thr Leu Thr Leu	2125	Asn Val Lys Thr Gln	2130	Glu Pro Asp Asn Leu	2135	Leu Phe Tyr Leu Gly	2140
Ser Thr Ala Ser	2145	Asp Phe Leu Ala Val	2150	Glu Met Arg Arg Gly	2155	Arg Val Ala Phe Leu	2160
	2165		2170		2175		2180

Asp Leu Gly Ser	Gly Ser Thr Arg Leu	Glu Phe Pro Asp Phe	Pro Ile Asp Asp Asn Arg	2185	2190	2195	2200
Trp His Ser Ile	His Val Ala Arg Phe	Gly Asn Ile Gly Ser	Leu Ser Val Lys Glu Met	2205	2210	2215	2220
Ser Ser Asn Gln	Lys Ser Pro Thr Lys	Thr Ser Lys Ser Pro	Gly Thr Ala Asn Val Leu	2225	2230	2235	2240
Asp Val Asn Asn	Ser Thr Leu Met Phe	Val Gly Gly Leu Gly	Gly Gln Ile Lys Lys Ser	2245	2250	2255	2260
Pro Ala Val Lys	Val Thr His Phe Lys	Gly Cys Leu Gly Glu	Ala Phe Leu Asn Gly Lys	2265	2270	2275	2280
Ser Ile Gly Leu	Trp Asn Tyr Ile Glu	Arg Glu Gly Lys Cys	Arg Gly Cys Phe Gly Ser	2285	2290	2295	2300
Ser Gln Asn Glu	Asp Pro Ser Phe His	Phe Asp Gly Ser Gly	Tyr Ser Val Val Glu Lys	2305	2310	2315	2320
Ser Leu Pro Ala	Thr Val Thr Gln Ile	Ile Met Leu Phe Asn	Thr Phe Ser Pro Asn Gly	2325	2330	2335	2340
Leu Leu Leu Tyr	Leu Gly Ser Tyr Gly	Thr Lys Asp Phe Leu	Ser Ile Glu Leu Phe Arg	2345	2350	2355	2360
Gly Arg Val Lys	Val Met Thr Asp Leu	Gly Ser Gly Pro Ile	Thr Leu Leu Thr Asp Arg	2365	2370	2375	2380
Arg Tyr Asn Asn	Gly Thr Trp Tyr Lys	Ile Ala Phe Gln Arg	Asn Arg Lys Gln Gly Val	2385	2390	2395	2400
Leu Ala Val Ile	Asp Ala Tyr Asn Thr	Ser Asn Lys Glu Thr	Lys Gln Gly Glu Thr Pro	2405	2410	2415	2420
Gly Ala Ser Ser	Asp Leu Asn Arg Leu	Asp Lys Asp Pro Ile	Tyr Val Gly Gly Leu Pro	2425	2430	2435	2440
Arg Ser Arg Val	Val Arg Arg Gly Val	Thr Thr Lys Ser Phe	Val Gly Cys Ile Lys Asn	2445	2450	2455	2460
Leu Glu Ile Ser	Arg Ser Thr Phe Asp	Leu Leu Arg Asn Ser	Tyr Gly Val Arg Lys Gly	2465	2470	2475	2480
Cys Leu Leu Glu	Pro Ile Arg Ser Val	Ser Phe Leu Lys Gly	Gly Tyr Ile Glu Leu Pro	2485	2490	2495	2500
Pro Lys Ser Leu	Ser Pro Glu Ser Glu	Trp Leu Val Thr Phe	Ala Thr Thr Asn Ser Ser	2505	2510	2515	2520
Gly Ile Ile Leu	Ala Ala Leu Gly Gly	Asp Val Glu Lys Arg	Gly Asp Arg Glu Glu Ala	2525	2530	2535	2540
His Val Pro Phe	Phe Ser Val Met Leu	Ile Gly Gly Asn Ile	Glu Val His Val Asn Pro	2545	2550	2555	2560
Gly Asp Gly Thr	Gly Leu Arg Lys Ala	Leu Leu His Ala Pro	Thr Gly Thr Cys Ser Asp	2565	2570	2575	2580
Gly Gln Ala His	Ser Ile Ser Leu Val	Arg Asn Arg Arg Ile	Ile Thr Val Gln Leu Asp	2585	2590	2595	2600
Glu Asn Asn Pro	Val Glu Met Lys Leu	Gly Thr Leu Val Glu	Ser Arg Thr Ile Asn Val	2605	2610	2615	2620
Ser Asn Leu Tyr	Val Gly Gly Ile Pro	Glu Gly Glu Gly Thr	Ser Leu Leu Thr Met Arg	2625	2630	2635	2640
Arg Ser Phe His	Gly Cys Ile Lys Asn	Leu Ile Phe Asn Leu	Glu Leu Leu Asp Phe Asn	2645	2650	2655	2660
Ser Ala Val Gly	His Glu Gln Val Asp	Leu Asp Thr Cys Trp	Leu Ser Glu Arg Pro Lys	2665	2670	2675	2680
Leu Ala Pro Asp	Ala Glu Asp Ser Lys	Leu Leu Arg Glu Pro	Arg Ala Phe Pro Glu Gln	2685	2690	2695	2700
Cys Val Val Asp	Ala Ala Leu Glu Tyr	Val Pro Gly Ala His	Gln Phe Gly Leu Thr Gln	2705	2710	2715	2720
Asn Ser His Phe	Ile Leu Pro Phe Asn	Gln Ser Ala Val Arg	Lys Lys Leu Ser Val Glu	2725	2730	2735	2740
Leu Ser Ile Arg	Thr Phe Ala Ser Ser	Gly Leu Ile Tyr Tyr	Met Ala His Gln Asn Gln	2745	2750	2755	2760
Ala Asp Tyr Ala	Val Leu Gln Leu His	Gly Gly Arg Leu His	Phe Met Phe Asp Leu Gly	2765	2770	2775	2780
Lys Gly Arg Thr	Lys Val Ser His Pro	Ala Leu Leu Ser Asp	Gly Lys Trp His Thr Val	2785	2790	2795	2800
Lys Thr Asp Tyr	Val Lys Arg Lys Gly	Phe Ile Thr Val Asp	Gly Arg Glu Ser Pro Met				

Val Thr Val Val	2805	2810	2815	2820
Gly Asp Gly Thr Met	2825	Leu Asp Val Glu Gly	Leu Phe Tyr Leu Gly	Gly
Leu Pro Ser Gln	2845	2830	2835	2840
Tyr Gln Ala Arg Lys	2850	Ile Gly Asn Ile Thr	His Ser Ile Pro Ala	Cys
Ile Gly Asp Val	2865	2870	2875	2880
Thr Val Asn Arg	2885	Gln Leu Asp Lys Asp	Ser Pro Val Ser Ala	Phe
Ala Ala Leu Val	2905	2890	2895	2900
Lys Glu Gly Tyr Lys	2910	Val Gln Ser Asp Val	Asn Ile Thr Leu Glu	Phe
Arg Thr Ser Ser	2925	2930	2935	2940
Gln Asn Gly Val Leu	2945	Leu Gly Ile Ser Thr	Ala Lys Val Asp Ala	Ile
Gly Leu Glu Leu	2965	2970	2975	2980
Val Asp Gly Lys Val	2985	Leu Phe His Val Asn	Asn Gly Ala Gly Arg	Ile
Thr Pro Ala Tyr	3005	3010	3015	3020
Glu Pro Lys Thr Ala	3025	3030	3035	3040
Thr Val Leu Cys Asp	3045	3050	3055	3060
Gln Ala Asn Lys	3065	3070		
Ser Lys His Arg Ile				
Thr Leu Ile Val Asp				
Val Asp Thr Asn Asn				
Pro Ile Tyr Val Gly				
Glu Ser Pro His				
Thr Gln Ser Thr Ser				
Val Lys Gln Lys Cys				
Leu Arg Ser Gln Thr				
Ser Phe Arg Gly Cys				
Arg Lys Leu Ala				
Leu Ile Lys Ser Pro				
Gln Val Gln Ser Phe				
Asp Phe Ser Arg Ala				
Glu Leu His Gly				
Val Phe Leu His Ser				
Cys Pro Gly Thr Glu				

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1786 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

- (D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P07942;

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Gly Leu Leu	Gln Leu Leu Ala Phe	Ser Phe Leu Ala Leu	Cys Arg Ala Arg Val Arg
1	5	10	15
Ala Gln Glu Pro	Glu Phe Ser Tyr Gly	Cys Ala Glu Gly Ser	Cys Tyr Pro Ala Thr Gly
25	30	35	40
Asp Leu Leu Ile	Gly Arg Ala Gln Lys	Leu Ser Val Thr Ser	Thr Cys Gly Leu His Lys
45	50	55	60
Pro Glu Pro Tyr	Cys Ile Val Ser His	Leu Gln Glu Asp Lys	Lys Cys Phe Ile Cys Asn
65	70	75	80
Ser Gln Asp Pro	Tyr His Glu Thr Leu	Asn Pro Asp Ser His	Leu Ile Glu Asn Val Val
85	90	95	100
Thr Thr Phe Ala	Pro Asn Arg Leu Lys	Ile Trp Trp Gln Ser	Glu Asn Gly Val Glu Asn
105	110	115	120
Val Thr Ile Gln	Leu Asp Leu Glu Ala	Glu Phe His Phe Thr	His Leu Ile Met Thr Phe
125	130	135	140
Lys Thr Phe Arg	Pro Ala Ala Met Leu	Ile Glu Arg Ser Ser	Asp Phe Gly Lys Thr Trp
145	150	155	160
Gly Val Tyr Arg	Tyr Phe Ala Tyr Asp	Cys Glu Ala Ser Phe	Pro Gly Ile Ser Thr Gly

Pro Met Lys Lys	165	170	175	180
Val Asp Asp Ile Ile	185	Cys Asp Ser Arg Tyr	Ser Asp Ile Glu Pro	Ser
Thr Glu Gly Glu	205	Leu Asp Pro Ala Phe	Lys Ile Glu Asp Pro	Tyr
Ser Pro Arg Ile	225	Ile Thr Asn Leu Arg	Ile Lys Phe Val Lys	Leu
His Thr Leu Gly	245	Ser Arg Met Glu Ile	Arg Glu Lys Tyr Tyr	Tyr
Ala Val Tyr Asp	265	Asn Cys Phe Cys Tyr	Gly His Ala Ser Glu	Cys
Ala Pro Val Asp	285	Val Glu Gly Met Val	His Gly His Cys Met	Cys
Arg His Asn Thr	305	Glu Leu Cys Met Asp	Phe Tyr His Asp Leu	Pro
Trp Arg Pro Ala	325	Asn Ala Cys Lys Lys	Cys Asn Cys Asn Glu	His
Ser Ile Ser Cys	345	Val Tyr Leu Ala Thr	Gly Asn Val Ser Gly	Gly
Val Cys Asp Asp	365	Met Gly Arg Asn Cys	Glu Gln Cys Lys Pro	Phe
Tyr Tyr Gln His	385	Arg Asp Pro Asn Phe	Cys Glu Arg Cys Thr	Cys
Asp Pro Ala Gly	405	Ile Cys Asp Ser Tyr	Thr Asp Phe Ser Thr	Gly
Leu Ile Ala Gly	425	Leu Asn Val Glu Gly	Glu His Cys Asp Val	Cys
Lys Glu Gly Phe	445	Glu Asp Pro Phe Gly	Cys Lys Ser Cys Ala	Cys
Asn Pro Leu Gly	465	Asn Pro Cys Asp Ser	Glu Thr Gly His Cys	Tyr
Cys Lys Arg Leu	485	Cys Asp Gln Cys Leu	Pro Glu His Trp Gly	Leu
Ser Asn Asp Leu	505	Cys Asp Cys Asp Leu	Gly Gly Ala Leu Asn	Asn
Ser Cys Phe Ala	525	Ser Cys Arg Pro His	Met Ile Gly Arg Gln	Cys
Asn Glu Val Glu	545	Ala Thr Leu Asp His	Tyr Leu Tyr Glu Ala	Glu
Glu Ala Asn Leu	565	Ile Val Glu Arg Gln	Tyr Ile Gln Asp Arg	Ile
Pro Ser Trp Thr	585	Arg Val Pro Glu Gly	Ala Tyr Leu Glu Phe	Phe
Ile Asp Asn Ile	605	Tyr Asp Ile Leu Ile	Arg Tyr Glu Pro Gln	Leu
Pro Asp His Trp	625	Thr Val Gln Arg Pro	Gly Arg Ile Pro Thr	Ser
Ser Arg Cys Gly	645	Asp Asp Asn Gln Val	Val Ser Leu Ser Pro	Gly
Ser Arg Tyr Val	665	Val Cys Phe Glu Lys	Gly Thr Asn Tyr Thr	Val
Arg Leu Glu Leu	685	Ser Asp Ser Asp Val	Glu Ser Pro Tyr Thr	Leu
Ile Asp Ser Leu	705	Cys Lys Ser Leu Asp	Ile Phe Thr Val Gly	Gly
Ser Gly Asp Gly	725	Ala Trp Glu Thr Phe	Gln Arg Tyr Arg Cys	Leu
Glu Asn Ser Arg	745	Pro Met Thr Asp Val	Cys Arg Asn Ile Ile	Phe
Ser Ile Ser Ala	765	Gly Leu Ala Cys Glu	Cys Asp Pro Gln Gly	Ser
Leu Ser Ser Val	785	Gly Gln Cys Gln Cys	Arg Pro Asn Val Val	Gly
				800

Arg	Thr	Cys	Asn	Arg	Cys	Ala	Pro	Gly	Thr	Phe	Gly	Phe	Gly	Pro	Ser	Gly	Cys	Lys	Pro
				805					810					815					820
Cys	Glu	Cys	His	Leu	Gln	Gly	Ser	Val	Asn	Ala	Phe	Cys	Asn	Pro	Val	Thr	Gly	Gln	Cys
				825					830					835					840
His	Cys	Phe	Gln	Gly	Val	Tyr	Ala	Arg	Gln	Cys	Asp	Arg	Cys	Leu	Pro	Gly	His	Trp	Gly
				845					850					855					860
Phe	Pro	Ser	Cys	Gln	Pro	Cys	Gln	Cys	Asn	Gly	His	Ala	Asp	Asp	Cys	Asp	Pro	Val	Thr
				865					870					875					880
Gly	Glu	Cys	Leu	Asn	Cys	Gln	Asp	Tyr	Thr	Met	Gly	His	Asn	Cys	Glu	Arg	Cys	Leu	Ala
				885					890					895					900
Gly	Tyr	Tyr	Gly	Asp	Pro	Ile	Ile	Gly	Ser	Gly	Asp	His	Cys	Arg	Pro	Cys	Pro	Cys	Pro
				905					910					915					920
Asp	Gly	Pro	Asp	Ser	Gly	Arg	Gln	Phe	Ala	Arg	Ser	Cys	Tyr	Gln	Asp	Pro	Val	Thr	Leu
				925					930					935					940
Gln	Leu	Ala	Cys	Val	Cys	Asp	Pro	Gly	Tyr	Ile	Gly	Ser	Arg	Cys	Asp	Asp	Cys	Ala	Ser
				945					950					955					960
Gly	Tyr	Phe	Gly	Asn	Pro	Ser	Glu	Val	Gly	Gly	Ser	Cys	Gln	Pro	Cys	Gln	Cys	His	Asn
				965					970					975					980
Asn	Ile	Asp	Thr	Thr	Asp	Pro	Glu	Ala	Cys	Asp	Lys	Glu	Thr	Gly	Arg	Cys	Leu	Lys	Cys
				985					990					995					1000
Leu	Tyr	His	Thr	Glu	Gly	Glu	His	Cys	Gln	Phe	Cys	Arg	Phe	Gly	Tyr	Tyr	Gly	Asp	Ala
				1005					1010					1015					1020
Leu	Arg	Gln	Asp	Cys	Arg	Lys	Cys	Val	Cys	Asn	Tyr	Leu	Gly	Thr	Val	Gln	Glu	His	Cys
				1025					1030					1035					1040
Asn	Gly	Ser	Asp	Cys	Gln	Cys	Asp	Lys	Ala	Thr	Gly	Gln	Cys	Leu	Cys	Leu	Pro	Asn	Val
				1045					1050					1055					1060
Ile	Gly	Gln	Asn	Cys	Asp	Arg	Cys	Ala	Pro	Asn	Thr	Trp	Gln	Leu	Ala	Ser	Gly	Thr	Gly
				1065					1070					1075					1080
Cys	Asp	Pro	Cys	Asn	Cys	Asn	Ala	Ala	His	Ser	Phe	Gly	Pro	Ser	Cys	Asn	Glu	Phe	Thr
				1085					1090					1095					1100
Gly	Gln	Cys	Gln	Cys	Met	Pro	Gly	Phe	Gly	Gly	Arg	Thr	Cys	Ser	Glu	Cys	Gln	Glu	Leu
				1105					1110					1115					1120
Phe	Trp	Gly	Asp	Pro	Asp	Val	Glu	Cys	Arg	Ala	Cys	Asp	Cys	Asp	Pro	Arg	Gly	Ile	Glu
				1125					1130					1135					1140
Thr	Pro	Gln	Cys	Asp	Gln	Ser	Thr	Gly	Gln	Cys	Val	Cys	Val	Glu	Gly	Val	Glu	Gly	Pro
				1145					1150										

Trp Gln Lys Ala	1425	Met Asp Leu Asp Gln	1430	Asp Val Leu Ser Ala	1435	Leu Ala Glu Val Glu Gln	1440
	1445	Val Ser Glu Ala Lys	1450	Leu Arg Ala Asp Glu	1455	Ala Lys Gln Ser Ala Glu	1460
Leu Ser Lys Met	1465	Lys Thr Asn Ala Thr	1470	Lys Glu Lys Met Asp	1475	Lys Ser Asn Glu Glu Leu	1480
Asp Ile Leu Leu	1485	Lys Gln Ile Arg Asn	1490	Phe Leu Thr Gln Asp	1495	Ser Ala Asp Leu Asp Ser	1500
Arg Asn Leu Ile	1505	Ala Asn Glu Val Leu	1510	Lys Met Glu Met Pro	1515	Ser Thr Pro Gln Gln Leu	1520
Ile Glu Ala Val	1525	Glu Asp Ile Arg Glu	1530	Arg Val Glu Ser Leu	1535	Ser Gln Val Glu Val Ile	1540
Gln Asn Leu Thr	1545	Ala Ala Asp Ile Ala	1550	Arg Ala Glu Met Leu	1555	Leu Glu Glu Ala Lys Arg	1560
Leu Gln His Ser	1565	Ala Thr Asp Val Lys	1570	Val Thr Ala Asp Met	1575	Val Lys Glu Ala Leu Glu	1580
Ala Ser Lys Ser	1585	Ala Gln Val Ala Ala	1590	Glu Lys Ala Ile Lys	1595	Gln Ala Asp Glu Asp Ile	1600
Glu Ala Glu Lys	1605	Asn Leu Leu Thr Ser	1610	Ile Glu Ser Glu Thr	1615	Ala Ala Ser Glu Glu Thr	1620
Gln Gly Thr Gln	1625	Ser Gln Arg Ile Ser	1630	Glu Leu Glu Arg Asn	1635	Val Glu Glu Leu Lys Arg	1640
Leu Phe Asn Ala	1645	Asn Ser Gly Glu Ala	1650	Glu Tyr Ile Glu Lys	1655	Val Val Tyr Thr Val Lys	1660
Lys Ala Ala Gln	1665	Asp Val Lys Lys Thr	1670	Leu Asp Gly Glu Leu	1675	Asp Glu Lys Tyr Lys Lys	1680
Gln Ser Ala Glu	1685	Ile Ala Lys Lys Thr	1690	Glu Glu Ser Ala Asp	1695	Ala Arg Arg Lys Ala Glu	1700
Val Glu Asn Leu	1705	Glu Ala Lys Thr Leu	1710	Leu Ala Gln Ala Asn	1715	Ser Lys Leu Gln Leu Leu	1720
Met Leu Gln Asn	1725	Arg Lys Tyr Glu Asp	1730	Asn Gln Arg Tyr Leu	1735	Glu Asp Lys Ala Gln Glu	1740
Lys Asp Leu Glu	1745	Glu Gly Glu Val Arg	1750	Ser Leu Leu Lys Asp	1755	Ile Ser Gln Lys Val Ala	1760
Leu Ala Arg Leu	1765	Cys Leu	1770		1775		1780
Val Tyr Ser Thr	1785						

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1786 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P02469

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Met Gly Leu Leu Gln Val Phe Ala Phe Gly Val Leu Ala Leu Trp Gly Thr Arg Val Cys	1	5	10	15	20
Ala Gln Glu Pro Glu Phe Ser Tyr Gly Cys Ala Glu Gly Ser Cys Tyr Pro Ala Thr Gly	25	30	35	40	45
Asp Leu Leu Ile Gly Arg Ala Gln Lys Leu Ser Val Thr Ser Thr Cys Gly Leu His Lys	50	55	60	65	70
Pro Glu Pro Tyr Cys Ile Val Ser His Leu Gln Glu Asp Lys Lys Cys Phe Ile Cys Asp	75	80	85	90	95

Ile	Asp	Ser	Leu	Val	Leu	Met	Pro	Tyr	Cys	Lys	Ser	Leu	Asp	Ile	Phe	Thr	Val	Gly	Gly	720
				705					710					715						
Ser	Gly	Asp	Gly	Glu	Val	Thr	Asn	Ser	Ala	Trp	Glu	Thr	Phe	Gln	Arg	Tyr	Arg	Cys	Leu	740
				725					730					735						
Glu	Asn	Ser	Arg	Ser	Val	Val	Lys	Thr	Pro	Met	Thr	Asp	Val	Cys	Arg	Asn	Ile	Ile	Phe	760
				745					750					755						
Ser	Ile	Ser	Ala	Leu	Ile	His	Gln	Thr	Gly	Leu	Ala	Cys	Glu	Cys	Asp	Pro	Gln	Gly	Ser	780
				765					770					775						
Leu	Ser	Ser	Val	Cys	Asp	Pro	Asn	Gly	Gly	Gln	Cys	Gln	Cys	Arg	Pro	Asn	Val	Val	Gly	800
				785					790					795						
Arg	Thr	Cys	Asn	Arg	Cys	Ala	Pro	Gly	Thr	Phe	Gly	Phe	Gly	Pro	Asn	Gly	Cys	Lys	Pro	820
				805					810					815						
Cys	Asp	Cys	His	Leu	Gln	Gly	Ser	Ala	Ser	Ala	Phe	Cys	Asp	Ala	Ile	Thr	Gly	Gln	Cys	840
				825					830					835						
His	Cys	Phe	Gln	Gly	Ile	Tyr	Ala	Arg	Gln	Cys	Asp	Arg	Cys	Leu	Pro	Gly	Tyr	Trp	Gly	860
				845					850					855						
Phe	Pro	Ser	Cys	Gln	Pro	Cys	Gln	Cys	Asn	Gly	His	Ala	Leu	Asp	Cys	Asp	Thr	Val	Thr	880
				865					870					875						
Gly	Glu	Cys	Leu	Ser	Cys	Gln	Asp	Tyr	Thr	Thr	Gly	His	Asn	Cys	Glu	Arg	Cys	Leu	Ala	900
				885					890					895						
Gly	Tyr	Tyr	Gly	Asp	Pro	Ile	Ile	Gly	Ser	Gly	Asp	His	Cys	Arg	Pro	Cys	Pro	Cys	Pro	920
				905					910					915						
Asp	Gly	Pro	Asp	Ser	Gly	Arg	Gln	Phe	Ala	Arg	Ser	Cys	Tyr	Gln	Asp	Pro	Val	Thr	Leu	940
				925					930					935						
Gln	Leu	Ala	Cys	Val	Cys	Asp	Pro	Gly	Tyr	Ile	Gly	Ser	Arg	Cys	Asp	Asp	Cys	Ala	Ser	960
				945					950					955						
Gly	Phe	Phe	Gly	Asn	Pro	Ser	Asp	Phe	Gly	Gly	Ser	Cys	Gln	Pro	Cys	Gln	Cys	His	His	980
				965					970					975						
Asn	Ile	Asp	Thr	Thr	Asp	Pro	Glu	Ala	Cys	Asp	Lys	Asp	Thr	Gly	Arg	Cys	Leu	Lys	Cys	1000
				985					990					995						
Leu	Tyr	His	Thr	Glu	Gly	Asp	His	Cys	Gln	Leu	Cys	Gln	Tyr	Gly	Tyr	Tyr	Gly	Asp	Ala	1020
				1005					1010					1015						
Leu	Arg	Gln	Asp	Cys	Arg	Lys	Cys	Val	Cys	Asn	Tyr	Leu	Gly	Thr	Val	Lys	Glu	His	Cys	1040
				1025					1030					1035						
Asn	Gly	Ser	Asp	Cys	His	Cys	Asp	Lys	Ala	Thr	Gly	Gln	Cys	Ser	Cys	Leu	Pro	Asn	Val	1060
				1045					1050					1055						
Ile	Gly	Gln	Asn	Cys	Asp	Arg	Cys	Ala	Pro	Asn	Thr	Trp	Gln	Leu	Ala	Ser	Gly	Thr	Gly	1080
				1065					1070					1075						
Cys	Gly	Pro	Cys	Asn	Cys	Asn	Ala	Ala	His	Ser	Phe	Gly	Pro	Ser	Cys	Asn	Glu	Phe	Thr	1100
				1085					1090					1095						
Gly	Gln	Cys	Gln	Cys	Met	Pro	Gly	Phe	Gly	Gly	Arg	Thr	Cys	Ser	Glu	Cys	Gln	Glu	Leu	1120
				1105					1110					1115						
Phe	Trp	Gly	Asp	Pro	Asp	Val	Glu	Cys	Arg	Ala	Cys	Asp	Cys	Asp	Pro	Arg	Gly	Ile	Glu	1140
				1125					1130					1135						
Thr	Pro	Gln	Cys	Asp	Gln	Ser	Thr	Gly	Gln	Cys	Val	Cys	Val	Glu	Gly	Val	Glu	Gly	Pro	1160
				1145					1150					1155						
Arg	Cys	Asp	Lys	Cys	Thr	Arg	Gly	Tyr	Ser	Gly	Val	Phe	Pro	Asp	Cys	Thr	Pro	Cys	His	1180
				1165					1170					1175						
Gln	Cys	Phe	Ala	Leu	Trp	Asp	Ala	Ile	Ile	Gly	Glu	Leu	Thr	Asn	Arg	Thr	His	Lys	Phe	1200
				1185					1190					1195						
Leu	Glu	Lys	Ala	Lys	Ala	Leu	Lys	Ile	Ser	Gly	Val	Ile	Gly	Pro	Tyr	Arg	Glu	Thr	Val	1220
				1205					1210					1215						
Asp	Ser	Val	Glu	Lys	Lys	Val	Asn	Glu	Ile	Lys	Asp	Ile	Leu	Ala	Gln	Ser	Pro	Ala	Ala	1240
				1225					1230					1235						
Glu	Pro	Leu	Lys	Asn	Ile	Gly	Ile	Leu	Phe	Glu	Glu	Ala	Glu	Lys	Leu	Thr	Lys	Asp	Val	1260
				1245					1250					1255						
Thr	Glu	Lys	Met	Ala	Gln	Val	Glu	Val	Lys	Leu	Thr	Asp	Thr	Ala	Ser	Gln	Ser	Asn	Ser	1280
				1265					1270					1275						
Thr	Ala	Gly	Glu	Leu	Gly	Ala	Leu	Gln	Ala	Glu	Ala	Glu	Ser	Leu	Asp	Lys	Thr	Val	Lys	1300
				1285					1290					1295						
Glu	Leu	Ala	Glu	Gln	Leu	Glu	Phe	Ile	Lys	Asn	Ser	Asp	Ile	Gln	Gly	Ala	Leu	Asp	Ser	1320
				1305					1310					1315						
Ile	Thr	Lys	Tyr	Phe	Gln	Met	Ser	Leu	Glu	Ala	Glu	Lys	Arg	Val	Asn	Ala	Ser	Thr	Thr	

Asp Pro Asn Ser	1325	1330	1335	1340
Thr Val Glu Gln Ser	1345	Ala Leu Thr Arg Asp	Arg Val Glu Asp Leu	Met
Leu Glu Arg Glu	1365	1350	1355	1360
Ser Pro Phe Lys Glu	1370	Gln Gln Glu Glu Gln	Ala Arg Leu Leu Asp	Glu
Leu Ala Gly Lys	1385	1390	1395	1400
Leu Gln Ser Leu Asp	1405	Leu Ser Ala Ala Ala	Gln Met Thr Cys Gly	Thr
Pro Pro Gly Ala	1425	1430	1435	1440
Asp Cys Ser Glu Ser	1445	Glu Cys Gly Gly Pro	Asn Cys Arg Thr Asp	Glu
Gly Glu Lys Lys	1465	1470	1475	1480
Cys Gly Gly Pro Gly	1485	Cys Gly Gly Leu Val	Thr Val Ala His Ser	Ala
Trp Gln Lys Ala	1505	1510	1515	1520
Met Asp Phe Asp Arg	1525	Asp Val Leu Ser Ala	Leu Ala Glu Val Glu	Gln
Leu Ser Lys Met	1545	1550	1555	1560
Val Ser Glu Ala Lys	1565	Val Arg Ala Asp Glu	Ala Lys Gln Asn Ala	Gln
Asp Val Leu Leu	1585	1590	1595	1600
Lys Thr Asn Ala Thr	1605	Lys Glu Lys Val Asp	Lys Ser Asn Glu Asp	Leu
Arg Asn Leu Ile	1625	1630	1635	1640
Lys Gln Ile Arg Asn	1645	Phe Leu Thr Glu Asp	Ser Ala Asp Leu Asp	Ser
Ile Glu Ala Val	1665	1670	1675	1680
Ala Asn Glu Val Leu	1685	Lys Ser Gly Asn Ala	Ser Thr Pro Gln Gln	Leu
Gln Asn Leu Thr	1705	1710	1715	1720
Glu Asp Ile Arg Glu	1725	Arg Val Glu Thr Leu	Ser Gln Val Glu Val	Ile
Leu Gln Gln Ser	1745	1750	1755	1760
Ala Ala Asp Ile Ala	1765	Arg Ala Glu Leu Leu	Leu Glu Glu Ala Lys	Arg
Ala Ser Lys Ser	1785	1790	1795	1800
Ala Thr Asp Val Lys		Val Thr Ala Asp Met	Val Lys Glu Ala Leu	Glu
Glu Ala Glu Lys		1805	1810	1815
Ala Gln Val Ala Ala		Glu Lys Ala Ile Lys	Gln Ala Asp Glu Asp	Ile
Gln Gly Thr Gln		1825	1830	1835
Asn Leu Leu Thr Ser		Ile Glu Ser Glu Thr	Ala Ala Ser Glu Glu	Thr
Leu Thr Asn Ala		1845	1850	1855
Ser Gln Arg Ile Ser		Lys Leu Glu Arg Asn	Val Glu Glu Leu Lys	Arg
Lys Ala Ala Gln		1865	1870	1875
Asn Ser Gly Glu Ala		Glu Tyr Ile Glu Lys	Val Val Tyr Ser Val	Lys
Gln Asn Ala Asp		1885	1890	1895
Asp Val Lys Lys Thr		Leu Asp Gly Glu Leu	Asp Glu Lys Tyr Lys	Lys
Val Glu Ser Leu		1905	1910	1915
Ile Ala Gln Lys Thr		Glu Glu Ser Ala Asp	Ala Arg Arg Lys Ala	Glu
Leu Leu Gln Asn		1925	1930	1935
Glu Ala Lys Thr Leu		Leu Ala Gln Ala Asn	Ser Lys Leu Gln Leu	Leu
Glu Asp Leu Glu		1945	1950	1955
Arg Lys Tyr Glu Asp		Asn Gln Lys Tyr Leu	Glu Asp Lys Ala Gln	Glu
Leu Val Arg Leu		1965	1970	1975
Glu Gly Glu Val Arg		Ser Leu Leu Lys Asp	Ile Ser Glu Lys Val	Ala
Val Tyr Ser Thr		1985		
Cys Leu				

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1801 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

- (D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P15800

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met	Glu	Trp	Ala	Ser	Gly	Lys	Pro	Gly	Arg	Gly	Arg	Gln	Gly	Gln	Pro	Val	Pro	Trp	Glu
1				5					10					15					20
Leu	Arg	Leu	Gly	Leu	Leu	Leu	Ser	Val	Leu	Ala	Ala	Thr	Leu	Ala	Gln	Val	Pro	Ser	Leu
				25					30					35					40
Asp	Val	Pro	Gly	Cys	Ser	Arg	Gly	Ser	Cys	Tyr	Pro	Ala	Thr	Gly	Asp	Leu	Leu	Val	Gly
				45					50					55					60
Arg	Ala	Asp	Arg	Leu	Thr	Ala	Ser	Ser	Thr	Cys	Gly	Leu	His	Ser	Pro	Gln	Pro	Tyr	Cys
				65					70					75					80
Ile	Val	Ser	His	Leu	Gln	Asp	Glu	Lys	Lys	Cys	Phe	Leu	Cys	Asp	Ser	Arg	Arg	Pro	Phe
				85					90					95					100
Ser	Ala	Arg	Asp	Asn	Pro	Asn	Ser	His	Arg	Ile	Gln	Asn	Val	Val	Thr	Ser	Phe	Ala	Pro
				105					110					115					120
Gln	Arg	Arg	Thr	Ala	Trp	Trp	Gln	Ser	Glu	Asn	Gly	Val	Pro	Met	Val	Thr	Ile	Gln	Leu
				125					130					135					140
Asp	Leu	Glu	Ala	Glu	Phe	His	Phe	Thr	His	Leu	Ile	Met	Thr	Phe	Lys	Thr	Phe	Arg	Pro
				145					150					155					160
Ala	Ala	Met	Leu	Val	Glu	Arg	Ser	Ala	Asp	Phe	Gly	Arg	Thr	Trp	Arg	Val	Tyr	Arg	Tyr
				165					170					175					180
Phe	Ser	Tyr	Asp	Cys	Gly	Ala	Asp	Phe	Pro	Gly	Ile	Pro	Leu	Ala	Pro	Pro	Arg	Arg	Trp
				185					190					195					200
Asp	Asp	Val	Val	Cys	Glu	Ser	Arg	Tyr	Ser	Glu	Ile	Glu	Pro	Ser	Thr	Glu	Gly	Glu	Val
				205					210					215					220
Ile	Tyr	Arg	Val	Leu	Asp	Pro	Ala	Ile	Pro	Ile	Pro	Asp	Pro	Tyr	Ser	Ser	Arg	Ile	Gln
				225					230					235					240
Asn	Leu	Leu	Lys	Ile	Thr	Asn	Leu	Arg	Val	Asn	Leu	Thr	Arg	Leu	His	Thr	Leu	Gly	Asp
				245					250					255					260
Asn	Leu	Leu	Asp	Pro	Arg	Arg	Glu	Ile	Arg	Glu	Lys	Tyr	Tyr	Tyr	Ala	Leu	Tyr	Glu	Leu
				265					270					275					280
Val	Ile	Arg	Gly	Asn	Cys	Phe	Cys	Tyr	Gly	His	Ala	Ser	Gln	Cys	Ala	Pro	Ala	Pro	Gly
				285					290					295					300
Ala	Pro	Ala	His	Ala	Glu	Gly	Met	Val	His	Gly	Ala	Cys	Ile	Cys	Lys	His	Asn	Thr	Arg
				305					310					315					320
Gly	Leu	Asn	Cys	Glu	Gln	Cys	Gln	Asp	Phe	Tyr	Gln	Asp	Leu	Pro	Trp	His	Pro	Ala	Glu
				325					330					335					340
Asp	Gly	His	Thr	His	Ala	Cys	Arg	Lys	Cys	Glu	Cys	Asn	Gly	His	Ser	His	Ser	Cys	His
				345					350					355					360
Phe	Asp	Met	Ala	Val	Tyr	Leu	Ala	Ser	Gly	Asn	Val	Ser	Gly	Gly	Val	Cys	Asp	Gly	Cys
				365					370					375					380
Gln	His	Asn	Thr	Ala	Gly	Arg	His	Cys	Glu	Leu	Cys	Arg	Pro	Phe	Phe	Tyr	Arg	Asp	Pro
				385					390					395					400
Thr	Lys	Asp	Met	Arg	Asp	Pro	Ala	Ala	Cys	Arg	Pro	Cys	Asp	Cys	Asp	Pro	Met	Gly	Ser
				405					410					415					420
Gln	Asp	Gly	Gly	Arg	Cys	Asp	Ser	His	Asp	Asp	Pro	Val	Leu	Gly	Leu	Val	Ser	Gly	Gln
				425					430					435					440
Cys	Arg	Cys	Lys	Glu	His	Val	Val	Gly	Thr	Arg	Cys	Gln	Gln	Cys	Arg	Asp	Gly	Phe	Phe
				445					450					455					460
Gly	Leu	Ser	Ala	Ser	Asn	Pro	Arg	Gly	Cys	Gln	Arg	Cys	Gln	Cys	Asn	Ser	Arg	Gly	Thr
				465					470					475					480
Val	Pro	Gly	Gly	Thr	Pro	Cys	Asp	Ser	Ser	Ser	Gly	Thr	Cys	Phe	Cys	Lys	Arg	Leu	Val
				485					490					495					500
Thr	Gly	Asp	Gly	Cys	Asp	Arg	Cys	Leu	Pro	Gly	His	Trp	Gly	Leu	Ser	His	Asp	Leu	Leu
				505					510					515					520
Gly	Cys	Arg	Pro	Cys	Asp	Cys	Asp	Val	Gly	Gly	Ala	Leu	Asp	Pro	Gln	Cys	Asp	Glu	Ala
				525					530					535					540
Thr	Gly	Gln	Cys	Pro	Cys	Arg	Pro	His	Met	Ile	Gly	Arg	Arg	Cys	Glu	Gln	Val	Gln	Pro
				545					550					555					560
Gly	Tyr	Phe	Arg	Pro	Phe	Leu	Asp	His	Leu	Thr	Trp	Glu	Ala	Glu	Gly	Ala	His	Gly	Gln
				565					570					575					580
Val	Leu	Glu	Val	Val	Glu	Arg	Leu	Val	Thr	Asn	Arg	Glu	Thr	Pro	Ser	Trp	Thr	Gly	Val
				585					590					595					600
Gly	Phe	Val	Arg	Leu	Arg	Glu	Gly	Gln	Glu	Val	Glu	Phe	Leu	Val	Thr	Ser	Leu	Pro	Arg

Ala Met Asp Tyr	605	Leu Leu Leu Arg	610	Trp Glu Pro Gln Val	615	Pro Glu Gln Trp Ala	620
Leu Glu Leu Val	625	Val Gln Arg Pro Gly	630	Pro Val Ser Ala His	635	Ser Pro Cys Gly His	640
Leu Pro Arg Asp	645	Asp Arg Ile Gln Gly	650	Met Leu His Pro Asn	655	Thr Arg Val Leu Val	660
Pro Arg Pro Val	665	Cys Leu Glu Pro Gly	670	Leu Ser Tyr Lys Leu	675	Lys Leu Lys Leu Thr	680
Thr Gly Gly Arg	685	Ala His Pro Glu Thr	690	Pro Tyr Ser Gly Ser	695	Gly Ile Leu Ile Asp	700
Leu Val Leu Gln	705	Pro His Val Leu Met	710	Leu Glu Met Phe Ser	715	Gly Gly Asp Ala Ala	720
Leu Glu Arg Arg	725	Thr Thr Phe Glu Arg	730	Tyr Arg Cys His Glu	735	Glu Gly Leu Met Pro	740
Lys Thr Pro Leu	745	Ser Glu Ala Cys Val	750	Pro Leu Leu Ile Ser	755	Ala Ser Ser Leu Val	760
Asn Gly Ala Leu	765	Pro Cys Gln Cys Asp	770	Pro Gln Gly Ser Leu	775	Ser Ser Glu Cys Asn	780
His Gly Gly Gln	785	Cys Arg Cys Lys Pro	790	Gly Val Val Gly Arg	795	Arg Cys Asp Ala Cys	800
Thr Gly Tyr Tyr	805	Gly Phe Gly Pro Ala	810	Gly Cys Gln Ala Cys	815	Gln Cys Ser Pro Asp	820
Ala Leu Ser Ala	825	Leu Cys Glu Gly Thr	830	Ser Gly Gln Cys Leu	835	Cys Arg Thr Gly Ala	840
Gly Leu Arg Cys	845	Asp His Cys Gln Arg	850	Gly Gln Trp Gly Phe	855	Pro Asn Cys Arg Pro	860
Val Cys Asn Gly	865	Arg Ala Asp Glu Cys	870	Asp Ala His Thr Gly	875	Ala Cys Leu Gly Cys	880
Asp Tyr Thr Gly	885	Gly Glu His Cys Glu	890	Arg Cys Ile Ala Gly	895	Phe His Gly Asp Pro	900
Leu Pro Tyr Gly	905	Gly Gln Cys Arg Pro	910	Cys Pro Cys Pro Glu	915	Gly Pro Gly Ser Gln	920
His Phe Ala Thr	925	Ser Cys His Arg Asp	930	Gly Tyr Ser Gln Gln	935	Ile Val Cys His Cys	940
Ala Gly Tyr Thr	945	Gly Leu Arg Cys Glu	950	Ala Cys Ala Pro Gly	955	His Phe Gly Asp Pro	960
Lys Pro Gly Gly	965	Arg Cys Gln Leu Cys	970	Glu Cys Ser Gly Asn	975	Ile Asp Pro Thr Asp	980
Gly Ala Cys Asp	985	Pro His Thr Gly Gln	990	Cys Leu Arg Cys Leu	995	His His Thr Glu Gly	1000
His Cys Gly His	1005	Cys Lys Pro Gly Phe	1010	His Gly Gln Ala Ala	1015	Arg Gln Ser Cys His	1020
Cys Thr Cys Asn	1025	Leu Leu Gly Thr Asp	1030	Pro Gln Arg Cys Pro	1035	Ser Thr Asp Leu Cys	1040
Cys Asp Pro Ser	1045	Thr Gly Gln Cys Pro	1050	Cys Leu Pro His Val	1055	Gln Gly Leu Ser Cys	1060
Arg Cys Ala Pro	1065	Asn Phe Trp Asn Phe	1070	Thr Ser Gly Arg Gly	1075	Cys Gln Pro Cys Ala	1080
His Pro Ser Arg	1085	Ala Arg Gly Pro Thr	1090	Cys Asn Glu Phe Thr	1095	Gly Gln Cys His Cys	1100
Ala Gly Phe Gly	1105	Gly Arg Thr Cys Ser	1110	Glu Cys Gln Glu Leu	1115	His Trp Gly Asp Pro	1120
Leu Gln Cys Arg	1125	Ala Cys Asp Cys Asp	1130	Pro Arg Gly Ile Asp	1135	Lys Pro Gln Cys His	1140
Ser Thr Gly His	1145	Cys Ser Cys Arg Pro	1150	Gly Val Ser Gly Val	1155	Arg Cys Asp Gln Cys	1160
Arg Gly Phe Ser	1165	Gly Val Phe Pro Ala	1170	Cys His Pro Cys His	1175	Ala Cys Phe Gly Asp	1180
Asp Arg Val Val	1185	Gln Asp Leu Ala Ala	1190	Arg Thr Arg Arg Leu	1195	Glu Gln Trp Ala Gln	1200
Leu Gln Gln Thr	1205	Gly Val Leu Gly Ala	1210	Phe Glu Ser Ser Phe	1215	Leu Asn Leu Gln Gly	1220
	1225		1230		1235		1240

Leu Gly Met Val	Gln Ala Ile Val Ala	Ala Arg Asn Thr Ser	Ala Ala Ser Thr Ala	Lys
1245	1250	1255	1260	
Leu Val Glu Ala	Thr Glu Gly Leu Arg	His Glu Ile Gly Lys	Thr Thr Glu Arg Leu	Thr
1265	1270	1275	1280	
Gln Leu Glu Ala	Glu Leu Thr Asp Val	Gln Asp Glu Asn Phe	Asn Ala Asn His Ala	Leu
1285	1290	1295	1300	
Ser Gly Leu Glu	Arg Asp Gly Leu Ala	Leu Asn Leu Thr Leu	Arg Gln Leu Asp Gln	His
1305	1310	1315	1320	
Leu Asp Ile Leu	Lys His Ser Asn Phe	Leu Gly Ala Tyr Asp	Ser Ile Arg His Ala	His
1325	1330	1335	1340	
Ser Gln Ser Thr	Glu Ala Glu Arg Arg	Ala Asn Ala Ser Thr	Phe Ala Ile Pro Ser	Pro
1345	1350	1355	1360	
Val Ser Asn Ser	Ala Asp Thr Arg Arg	Arg Ala Glu Val Leu	Met Gly Ala Gln Arg	Glu
1365	1370	1375	1380	
Asn Phe Asn Arg	Gln His Leu Ala Asn	Gln Gln Ala Leu Gly	Arg Leu Ser Thr His	Thr
1385	1390	1395	1400	
His Thr Leu Ser	Leu Thr Gly Val Asn	Glu Leu Val Cys Gly	Ala Pro Gly Asp Ala	Pro
1405	1410	1415	1420	
Cys Ala Thr Ser	Pro Cys Gly Gly Ala	Gly Cys Arg Asp Glu	Asp Gly Gln Pro Arg	Cys
1425	1430	1435	1440	
Gly Gly Leu Gly	Cys Ser Gly Ala Ala	Ala Thr Ala Asp Leu	Ala Leu Gly Arg Ala	Arg
1445	1450	1455	1460	
His Thr Gln Ala	Glu Leu Gln Arg Ala	Leu Val Glu Gly Gly	Gly Ile Leu Ser Arg	Val
1465	1470	1475	1480	
Ser Glu Thr Arg	Arg Gln Ala Glu Glu	Ala Gln Gln Arg Ala	Gln Ala Ala Leu Asp	Lys
1485	1490	1495	1500	
Ala Asn Ala Ser	Arg Gly Gln Val Glu	Gln Ala Asn Gln Glu	Leu Arg Glu Leu Ile	Gln
1505	1510	1515	1520	
Asn Val Lys Asp	Phe Leu Ser Gln Glu	Gly Ala Asp Pro Asp	Ser Ile Glu Met Val	Ala
1525	1530	1535	1540	
Thr Arg Val Leu	Asp Ile Ser Ile Pro	Ala Ser Pro Glu Gln	Ile Gln Arg Leu Ala	Ser
1545	1550	1555	1560	
Glu Ile Ala Glu	Arg Val Arg Ser Leu	Ala Asp Val Asp Thr	Ile Leu Ala His Thr	Met
1565	1570	1575	1580	
Gly Asp Val Arg	Arg Ala Glu Gln Leu	Leu Gln Asp Ala Gln	Arg Ala Arg Ser Arg	Ala
1585	1590	1595	1600	
Glu Gly Glu Arg	Gln Lys Ala Glu Thr	Val Gln Ala Ala Leu	Glu Glu Ala Gln Arg	Ala
1605	1610	1615	1620	
Gln Gly Ala Ala	Gln Gly Ala Ile Arg	Gly Ala Val Val Asp	Thr Lys Asn Thr Glu	Gln
1625	1630	1635	1640	
Thr Leu Gln Gln	Val Gln Glu Arg Met	Ala Gly Thr Glu Gln	Ser Leu Asn Ser Ala	Ser
1645	1650	1655	1660	
Glu Arg Ala Arg	Gln Leu His Ala Leu	Leu Glu Ala Leu Lys	Leu Lys Arg Ala Gly	Asn
1665	1670	1675	1680	
Ser Leu Ala Ala	Ser Thr Ala Glu Glu	Thr Ala Gly Ser Ala	Gln Ser Arg Ala Arg	Glu
1685	1690	1695	1700	
Ala Glu Lys Gln	Leu Arg Glu Gln Val	Gly Asp Gln Tyr Gln	Thr Val Arg Ala Leu	Ala
1705	1710	1715	1720	
Glu Arg Lys Ala	Glu Gly Val Leu Ala	Ala Gln Ala Arg Ala	Glu Gln Leu Arg Asp	Glu
1725	1730	1735	1740	
Ala Arg Gly Leu	Leu Gln Ala Ala Gln	Asp Lys Leu Gln Arg	Leu Gln Glu Leu Glu	Gly
1745	1750	1755	1760	
Thr Tyr Glu Glu	Asn Glu Arg Glu Leu	Glu Val Lys Ala Ala	Gln Leu Asp Gly Leu	Glu
1765	1770	1775	1780	
Ala Arg Met Arg	Ser Val Leu Gln Ala	Ile Asn Leu Gln Val	Gln Ile Tyr Asn Thr	Cys
1785	1790	1795	1800	
Gln				

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1798 AMINO ACIDS
 (B) TYPE: AMINO ACID
 (C) STRANDEDNESS:
 (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF
 GENE BANK ACCESSION NUMBER P55268

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Met	Glu	Leu	Thr	Ser	Arg	Glu	Arg	Gly	Arg	Gly	Gln	Pro	Leu	Pro	Trp	Glu	Leu	Arg	Leu	1	5	10	15	20
Gly	Leu	Leu	Leu	Ser	Val	Leu	Ala	Ala	Thr	Leu	Ala	Gln	Ala	Pro	Ala	Pro	Asp	Val	Pro	25	30	35	40	45
Gly	Cys	Ser	Arg	Gly	Ser	Cys	Tyr	Pro	Ala	Thr	Gly	Asp	Leu	Leu	Val	Gly	Arg	Ala	Asp	50	55	60	65	70
Arg	Leu	Thr	Ala	Ser	Ser	Thr	Cys	Gly	Leu	Asn	Gly	Pro	Gln	Pro	Tyr	Cys	Ile	Val	Ser	75	80	85	90	95
His	Leu	Gln	Asp	Glu	Lys	Lys	Cys	Phe	Leu	Cys	Asp	Ser	Arg	Arg	Pro	Phe	Ser	Ala	Arg	100	105	110	115	120
Asp	Asn	Pro	His	Ser	His	Arg	Ile	Gln	Asn	Val	Val	Thr	Ser	Phe	Ala	Pro	Gln	Arg	Arg	125	130	135	140	145
Ala	Ala	Trp	Trp	Gln	Ser	Glu	Asn	Gly	Ile	Pro	Ala	Val	Thr	Ile	Gln	Leu	Asp	Leu	Glu	150	155	160	165	170
Ala	Glu	Phe	His	Phe	Thr	His	Leu	Ile	Met	Thr	Phe	Lys	Thr	Phe	Arg	Pro	Ala	Ala	Met	175	180	185	190	195
Leu	Val	Glu	Arg	Ser	Ala	Asp	Phe	Gly	Arg	Thr	Trp	His	Val	Tyr	Arg	Tyr	Phe	Ser	Tyr	200	205	210	215	220
Asp	Cys	Gly	Ala	Asp	Phe	Pro	Gly	Val	Pro	Leu	Ala	Pro	Pro	Arg	His	Trp	Asp	Asp	Val	225	230	235	240	245
Val	Cys	Glu	Ser	Arg	Tyr	Ser	Glu	Ile	Glu	Pro	Ser	Thr	Glu	Gly	Glu	Val	Ile	Tyr	Arg	250	255	260	265	270
Val	Leu	Asp	Pro	Ala	Ile	Pro	Ile	Pro	Asp	Pro	Tyr	Ser	Ser	Arg	Ile	Gln	Asn	Leu	Leu	275	280	285	290	295
Lys	Ile	Thr	Asn	Leu	Arg	Val	Asn	Leu	Thr	Arg	Leu	His	Thr	Leu	Gly	Asp	Asn	Leu	Leu	300	305	310	315	320
Asp	Pro	Arg	Arg	Glu	Ile	Arg	Glu	Lys	Tyr	Tyr	Tyr	Ala	Leu	Tyr	Glu	Leu	Val	Val	Arg	325	330	335	340	345
Gly	Asn	Cys	Phe	Cys	Tyr	Gly	His	Ala	Ser	Glu	Cys	Ala	Pro	Ala	Pro	Gly	Ala	Pro	Ala	350	355	360	365	370
His	Ala	Glu	Gly	Met	Val	His	Gly	Ala	Cys	Ile	Cys	Lys	His	Asn	Thr	Arg	Gly	Leu	Asn	375	380	385	390	395
Cys	Glu	Gln	Cys	Gln	Asp	Phe	Tyr	Arg	Asp	Leu	Pro	Trp	Arg	Pro	Ala	Glu	Asp	Gly	His	400	405	410	415	420
Ser	His	Ala	Cys	Arg	Lys	Cys	Glu	Cys	His	Gly	His	Thr	His	Ser	Cys	His	Phe	Asp	Met	425	430	435	440	445
Ala	Val	Tyr	Leu	Ala	Ser	Gly	Asn	Val	Ser	Gly	Gly	Val	Cys	Asp	Gly	Cys	Gln	His	Asn	450	455	460	465	470
Thr	Ala	Gly	Arg	His	Cys	Glu	Leu	Cys	Arg	Pro	Phe	Phe	Tyr	Arg	Asp	Pro	Thr	Lys	Asp	475	480	485	490	495
Leu	Arg	Asp	Pro	Ala	Val	Cys	Arg	Ser	Cys	Asp	Cys	Asp	Pro	Met	Gly	Ser	Gln	Asp	Gly					
Gly	Arg	Cys	Asp	Ser	His	Asp	Asp	Pro	Ala	Leu	Gly	Leu	Val	Ser	Gly	Gln	Cys	Arg	Cys					
Lys	Glu	His	Val	Val	Gly	Thr	Arg	Cys	Gln	Gln	Cys	Arg	Asp	Gly	Phe	Phe	Gly	Leu	Ser					
Ile	Ser	Asp	Arg	Leu	Gly	Cys	Arg	Arg	Cys	Gln	Cys	Asn	Ala	Arg	Gly	Thr	Val	Pro	Gly					
Ser	Thr	Pro	Cys	Asp	Pro	Asn	Ser	Gly	Ser	Cys	Tyr	Cys	Lys	Arg	Leu	Val	Thr	Gly	Arg					

Gly	Cys	Asp	Arg	Cys	Leu	Pro	Gly	His	Trp	Gly	Leu	Ser	His	Asp	Leu	Leu	Gly	Cys	Arg
				505					510					515					520
Pro	Cys	Asp	Cys	Asp	Val	Gly	Gly	Ala	Leu	Asp	Pro	Gln	Cys	Asp	Glu	Gly	Thr	Gly	Gln
				525					530					535					540
Cys	His	Cys	Arg	Gln	His	Met	Val	Gly	Arg	Arg	Cys	Glu	Gln	Val	Gln	Pro	Gly	Tyr	Phe
				545					550					555					560
Arg	Pro	Phe	Leu	Asp	His	Leu	Ile	Trp	Glu	Ala	Glu	Asp	Thr	Arg	Gly	Gln	Val	Leu	Asp
				565					570					575					580
Val	Val	Glu	Arg	Leu	Val	Thr	Pro	Gly	Glu	Thr	Pro	Ser	Trp	Thr	Gly	Ser	Gly	Phe	Val
				585					590					595					600
Arg	Leu	Gln	Glu	Gly	Gln	Thr	Leu	Glu	Phe	Leu	Val	Ala	Ser	Val	Pro	Lys	Ala	Met	Asp
				605					610					615					620
Tyr	Asp	Leu	Leu	Leu	Arg	Leu	Glu	Pro	Gln	Val	Pro	Glu	Gln	Trp	Ala	Glu	Leu	Glu	Leu
				625					630					635					640
Ile	Val	Gln	Arg	Pro	Gly	Pro	Val	Pro	Ala	His	Ser	Leu	Cys	Gly	His	Leu	Val	Pro	Lys
				645					650					655					660
Asp	Asp	Arg	Ile	Gln	Gly	Thr	Leu	Gln	Pro	His	Ala	Arg	Tyr	Leu	Ile	Phe	Pro	Asn	Pro
				665					670					675					680
Val	Cys	Leu	Glu	Pro	Gly	Ile	Ser	Tyr	Lys	Leu	His	Leu	Lys	Leu	Val	Arg	Thr	Gly	Gly
				685					690					695					700
Ser	Ala	Gln	Pro	Glu	Thr	Pro	Tyr	Ser	Gly	Pro	Gly	Leu	Leu	Ile	Asp	Ser	Leu	Val	Leu
				705					710					715					720
Leu	Pro	Arg	Val	Leu	Val	Leu	Glu	Met	Phe	Ser	Gly	Gly	Asp	Ala	Ala	Ala	Leu	Glu	Arg
				725					730					735					740
Gln	Ala	Thr	Phe	Glu	Arg	Tyr	Gln	Cys	His	Glu	Glu	Gly	Leu	Val	Pro	Ser	Lys	Thr	Ser
				745					750					755					760
Pro	Ser	Glu	Ala	Cys	Ala	Pro	Leu	Leu	Ile	Ser	Leu	Ser	Thr	Leu	Ile	Tyr	Asn	Gly	Ala
				765					770					775					780
Leu	Pro	Cys	Gln	Cys	Asn	Pro	Gln	Gly	Ser	Leu	Ser	Ser	Glu	Cys	Asn	Pro	His	Gly	Gly
				785					790					795					800
Gln	Cys	Leu	Cys	Lys	Pro	Gly	Val	Val	Gly	Arg	Arg	Cys	Asp	Leu	Cys	Ala	Pro	Gly	Tyr
				805					810					815					820
Tyr	Gly	Phe	Gly	Pro	Thr	Gly	Cys	Gln	Ala	Cys	Gln	Cys	Ser	His	Glu	Gly	Ala	Leu	Ser
				825					830					835					840
Ser	Leu	Cys	Glu	Lys	Thr	Ser	Gly	Gln	Cys	Leu	Cys	Arg	Thr	Gly	Ala	Phe	Gly	Leu	Arg
				845					850					855					860
Cys	Asp	Arg	Cys	Gln	Arg	Gly	Gln	Trp	Gly	Phe	Pro	Ser	Cys	Arg	Pro	Cys	Val	Cys	Asn
				865					870					875					880
Gly	His	Ala	Asp	Glu	Cys	Asn	Thr	His	Thr	Gly	Ala	Cys	Leu	Gly	Cys	Arg	Asp	His	Thr
				885					890					895					900
Gly	Gly	Glu	His	Cys	Glu	Arg	Cys	Ile	Ala	Gly	Phe	His	Arg	Asp	Pro	Arg	Leu	Pro	Tyr
				905					910					915					920
Gly	Gly	Gln	Cys	Arg	Pro	Cys	Pro	Cys	Pro	Glu	Gly	Pro	Gly	Ser	Gln	Arg	His	Phe	Ala
				925					930					935					940
Thr	Ser	Cys	His	Gln	Asp	Glu	Tyr	Ser	Gln	Gln	Ile	Val	Cys	His	Cys	Arg	Ala	Gly	Tyr
				945					950					955					960
Thr	Gly	Leu	Arg	Cys	Glu	Ala	Cys	Ala	Pro	Gly	His	Phe	Gly	Asp	Pro	Ser	Arg	Pro	Gly
				965					970					975					980
Gly	Arg	Cys	Gln	Leu	Cys	Glu	Cys	Ser	Gly	Asn	Ile	Asp	Pro	Met	Asp	Pro	Asp	Ala	Cys
				985					990					995					1000
Asp	Pro	His	Thr	Gly	Gln	Cys	Leu	Arg	Cys	Leu	His	His	Thr	Glu	Gly	Pro	His	Cys	Ala
				1005					1010					1015					1020
His	Cys	Lys	Pro	Gly	Phe	His	Gly	Gln	Ala	Ala	Arg	Gln	Ser	Cys	His	Arg	Cys	Thr	Cys
				1025					1030					1035					1040
Asn	Leu	Leu	Gly	Thr	Asn	Pro	Gln	Gln	Cys	Pro	Ser	Pro	Asp	Gln	Cys	His	Cys	Asp	Pro
				1045					1050					1055					1060
Ser	Ser	Gly	Gln	Cys	Pro	Cys	Leu	Pro	Asn	Val	Gln	Gly	Pro	Ser	Cys	Asp	Arg	Cys	Ala
				1065					1070					1075					1080
Pro	Asn	Phe	Trp	Asn	Leu	Thr	Ser	Gly	His	Gly	Cys	Gln	Pro	Cys	Ala	Cys	His	Pro	Ser
				1085					1090					1095					1100
Arg	Ala	Arg	Gly	Pro	Thr	Cys	Asn	Glu	Phe	Thr	Gly	Gln	Cys	His	Cys	Arg	Ala	Gly	Phe
				1105					1110					1115					1120
Gly	Gly	Arg	Thr	Cys	Ser	Glu	Cys	Gln	Glu	Leu	His	Trp	Gly	Asp	Pro	Gly	Leu	Gln	Cys

His Ala Cys Asp	1125	Cys Asp Ser Arg Gly	1130	Ile Asp Thr Pro Gln	1135	Cys His Arg Phe Thr Gly	1140
His Cys Ser Cys	1145	Arg Pro Gly Val Ser	1150	Gly Val Arg Cys Asp	1155	Gln Cys Ala Arg Gly Phe	1160
Ser Gly Ile Phe	1165	Pro Ala Cys His Pro	1170	Cys His Ala Cys Phe	1175	Gly Asp Trp Asp Arg Val	1180
Val Gln Asp Leu	1185	Ala Ala Arg Thr Gln	1190	Arg Leu Glu Gln Arg	1195	Ala Gln Glu Leu Gln Gln	1200
Thr Gly Val Leu	1205	Gly Ala Phe Glu Ser	1210	Ser Phe Trp His Met	1215	Gln Glu Lys Leu Gly Ile	1220
Val Gln Gly Ile	1225	Val Gly Ala Arg Asn	1230	Thr Ser Ala Ala Ser	1235	Thr Ala Gln Leu Val Glu	1240
Ala Thr Glu Glu	1245	Leu Arg Arg Glu Ile	1250	Gly Glu Ala Thr Glu	1255	His Leu Thr Gln Leu Glu	1260
Ala Asp Leu Thr	1265	Asp Val Gln Asp Glu	1270	Asn Phe Asn Ala Asn	1275	His Ala Leu Ser Gly Leu	1280
Glu Arg Asp Arg	1285	Leu Ala Leu Asn Leu	1290	Thr Leu Arg Gln Leu	1295	Asp Gln His Leu Asp Leu	1300
Leu Lys His Ser	1305	Asn Phe Leu Gly Ala	1310	Tyr Asp Ser Ile Arg	1315	His Ala His Ser Gln Ser	1320
Ala Glu Ala Glu	1325	Arg Arg Ala Asn Thr	1330	Ser Ala Leu Ala Val	1335	Pro Ser Pro Val Ser Asn	1340
Ser Ala Ser Ala	1345	Arg His Arg Thr Glu	1350	Ala Leu Met Asp Ala	1355	Gln Lys Glu Asp Phe Asn	1360
Ser Lys His Met	1365	Ala Asn Gln Arg Ala	1370	Leu Gly Lys Leu Ser	1375	Ala His Thr His Thr Leu	1380
Ser Leu Thr Asp	1385	Ile Asn Glu Leu Val	1390	Cys Gly Ala Pro Gly	1395	Asp Ala Pro Cys Ala Thr	1400
Ser Pro Cys Gly	1405	Gly Ala Gly Cys Arg	1410	Asp Glu Asp Gly Gln	1415	Pro Arg Cys Gly Gly Leu	1420
Ser Cys Asn Gly	1425	Ala Ala Ala Thr Ala	1430	Asp Leu Ala Leu Gly	1435	Arg Ala Arg His Thr Gln	1440
Ala Glu Leu Gln	1445	Arg Ala Leu Ala Glu	1450	Gly Gly Ser Ile Leu	1455	Ser Arg Val Ala Glu Thr	1460
Arg Arg Gln Ala	1465	Ser Glu Ala Gln Gln	1470	Arg Ala Gln Ala Ala	1475	Leu Asp Lys Ala Asn Ala	1480
Ser Arg Gly Gln	1485	Val Glu Gln Ala Asn	1490	Gln Glu Leu Gln Glu	1495	Leu Ile Gln Ser Val Lys	1500
Asp Phe Leu Asn	1505	Gln Glu Gly Ala Asp	1510	Pro Asp Ser Ile Glu	1515	Met Val Ala Thr Arg Val	1520
Leu Glu Leu Ser	1525	Ile Pro Ala Ser Ala	1530	Glu Gln Ile Gln His	1535	Leu Ala Gly Ala Ile Ala	1540
Glu Arg Val Arg	1545	Ser Leu Ala Asp Val	1550	Asp Ala Ile Leu Ala	1555	Arg Thr Val Gly Asp Val	1560
Arg Arg Ala Glu	1565	Gln Leu Leu Gln Asp	1570	Ala Arg Arg Ala Arg	1575	Ser Trp Ala Glu Asp Glu	1580
Lys Gln Lys Ala	1585	Glu Thr Val Gln Ala	1590	Ala Leu Glu Glu Ala	1595	Gln Arg Ala Gln Gly Ile	1600
Ala Gln Gly Ala	1605	Ile Arg Gly Ala Val	1610	Ala Asp Thr Arg Asp	1615	Thr Glu Gln Thr Leu Tyr	1620
Gln Val Gln Glu	1625	Arg Met Ala Gly Ala	1630	Glu Arg Ala Leu Ser	1635	Ser Ala Gly Glu Arg Ala	1640
Arg Gln Leu Asp	1645	Ala Leu Leu Glu Ala	1650	Leu Lys Leu Lys Arg	1655	Ala Gly Asn Ser Leu Ala	1660
Ala Ser Thr Ala	1665	Glu Glu Thr Ala Gly	1670	Ser Ala Gln Gly Arg	1675	Ala Gln Glu Ala Glu Gln	1680
Leu Leu Arg Gly	1685	Pro Leu Gly Asp Gln	1690	Tyr Gln Thr Val Lys	1695	Ala Leu Ala Glu Arg Lys	1700
Ala Gln Gly Val	1705	Leu Ala Ala Gln Ala	1710	Arg Ala Glu Gln Leu	1715	Arg Asp Glu Ala Arg Asp	1720
Leu Leu Gln Ala	1725	Ala Gln Asp Lys Leu	1730	Gln Arg Leu Gln Glu	1735	Leu Glu Gly Thr Tyr Glu	1740
	1745		1750		1755		1760

Glu Asn Glu Arg	Ala Leu Glu Ser Lys	Ala Ala Gln Leu Asp	Gly Leu Glu Ala Arg	Met
	1765	1770	1775	1780
Arg Ser Val Leu	Gln Ala Ile Asn Leu	Gln Val Gln Ile Tyr	Asn Thr Cys Gln	
	1785	1790	1795	

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1607 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P02468

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Thr Gly Gly	Gly Arg Ala Ala	Leu Ala Leu Gln Pro	Arg Gly Arg Leu Trp	Pro Leu
1	5	10	15	20
Leu Ala Val Leu	Ala Ala Val Ala	Gly Cys Val Arg Ala	Ala Met Asp Glu Cys	Ala Asp
	25	30	35	40
Glu Gly Gly Arg	Pro Gln Arg Cys Met	Pro Glu Phe Val Asn	Ala Ala Phe Asn Val	Thr
	45	50	55	60
Val Val Ala Thr	Asn Thr Cys Gly Thr	Pro Pro Glu Glu Tyr	Cys Val Gln Thr Gly	Val
	65	70	75	80
Thr Gly Val Thr	Lys Ser Cys His Leu	Cys Asp Ala Gly Gln	Gln His Leu Gln His	Gly
	85	90	95	100
Ala Ala Phe Leu	Thr Asp Tyr Asn Asn	Gln Ala Asp Thr Thr	Trp Trp Gln Ser Gln	Thr
	105	110	115	120
Met Leu Ala Gly	Val Gln Tyr Pro Asn	Ser Ile Asn Leu Thr	Leu His Leu Gly Lys	Ala
	125	130	135	140
Phe Asp Ile Thr	Tyr Val Arg Leu Lys	Phe His Thr Ser Arg	Pro Glu Ser Phe Ala	Ile
	145	150	155	160
Tyr Lys Arg Thr	Arg Glu Asp Gly Pro	Trp Ile Pro Tyr Gln	Tyr Tyr Ser Gly Ser	Cys
	165	170	175	180
Glu Asn Thr Tyr	Ser Lys Ala Asn Arg	Gly Phe Ile Arg Thr	Gly Gly Asp Glu Gln	Gln
	185	190	195	200
Ala Leu Cys Thr	Asp Glu Phe Ser Asp	Ile Ser Pro Leu Thr	Gly Gly Asn Val Ala	Phe
	205	210	215	220
Ser Thr Leu Glu	Gly Arg Pro Ser Ala	Tyr Asn Phe Asp Asn	Ser Pro Val Leu Gln	Glu
	225	230	235	240
Trp Val Thr Ala	Thr Asp Ile Arg Val	Thr Leu Asn Arg Leu	Asn Thr Phe Gly Asp	Glu
	245	250	255	260
Val Phe Asn Glu	Pro Lys Val Leu Lys	Ser Tyr Tyr Tyr Ala	Ile Ser Asp Phe Ala	Val
	265	270	275	280
Gly Gly Arg Cys	Lys Cys Asn Gly His	Ala Ser Glu Cys Val	Lys Asn Glu Phe Asp	Lys
	285	290	295	300
Leu Met Cys Asn	Cys Lys His Asn Thr	Tyr Gly Val Asp Cys	Glu Lys Cys Leu Pro	Phe
	305	310	315	320
Phe Asn Asp Arg	Pro Trp Arg Arg Ala	Thr Ala Glu Ser Ala	Ser Glu Ser Leu Pro	Cys
	325	330	335	340
Asp Cys Asn Gly	Arg Ser Gln Glu Cys	Tyr Phe Asp Pro Glu	Leu Tyr Arg Ser Thr	Gly
	345	350	355	360
His Gly Gly His	Cys Thr Asn Cys Arg	Asp Asn Thr Asp Gly	Ala Lys Cys Glu Arg	Cys
	365	370	375	380
Arg Glu Asn Phe	Phe Arg Leu Gly Asn	Thr Glu Ala Cys Ser	Pro Cys His Cys Ser	Pro
	385	390	395	400
Val Gly Ser Leu	Ser Thr Gln Cys Asp	Ser Tyr Gly Arg Cys	Ser Cys Lys Pro Gly	Val

Met Gly Asp Lys	405	410	415	420
Cys Asp Arg Cys Gln	425	430	435	440
Arg Pro Cys Ser	445	450	455	460
Cys Val Cys Lys	465	470	475	480
Asn Leu Glu Ser	485	490	495	500
Val Cys Thr Asn	505	510	515	520
Glu Asp Gly Trp	525	530	535	540
Asp Arg Gln Asp	545	550	555	560
Val Lys Phe Leu	565	570	575	580
Val Asp Arg Arg	585	590	595	600
Arg Val Ser Val	605	610	615	620
Tyr Ile Phe Arg	625	630	635	640
Glu Phe Gln Lys	645	650	655	660
Arg Thr Ala Gly	665	670	675	680
Pro Ala Thr Trp	685	690	695	700
Thr Cys Leu Pro	705	710	715	720
Leu Cys Thr Cys	725	730	735	740
Arg Asp Asn Thr	745	750	755	760
Thr Leu Gly Thr	765	770	775	780
Ile Val Pro Lys	785	790	795	800
Arg Cys Glu Leu	805	810	815	820
Arg Leu Cys Arg	825	830	835	840
Asn Arg Leu Thr	845	850	855	860
Arg Cys Lys Glu	865	870	875	880
Ala Cys Ala Cys	885	890	895	900
Gly Gln Cys Gln	905	910	915	920
Tyr Tyr Asn Leu	925	930	935	940
Thr Asn Gly Gln	945	950	955	960
Gln His Cys Glu	965	970	975	980
Cys Asp Cys His	985	990	995	1000
Cys Arg Glu Gly	1005	1010	1015	1020
Arg Ser Trp Pro	1025	1030	1035	1040

Ala Glu His Arg	Val Lys Leu Gln Glu	Leu Glu Ser Leu Ile	Ala Asn Leu Gly Thr Gly	1045	1050	1055	1060
Asp Asp Met Val	Thr Asp Gln Ala Phe	Glu Asp Arg Leu Lys	Glu Ala Glu Arg Glu Val	1065	1070	1075	1080
Thr Asp Leu Leu	Arg Glu Ala Gln Glu	Val Lys Asp Val Asp	Gln Asn Leu Met Asp Arg	1085	1090	1095	1100
Leu Gln Arg Val	Asn Ser Ser Leu His	Ser Gln Ile Ser Arg	Leu Gln Asn Ile Arg Asn	1105	1110	1115	1120
Thr Ile Glu Glu	Thr Gly Ile Leu Ala	Glu Arg Ala Arg Ser	Arg Val Glu Ser Thr Glu	1125	1130	1135	1140
Gln Leu Ile Glu	Ile Ala Ser Arg Glu	Leu Glu Lys Ala Lys	Met Ala Ala Ala Asn Val	1145	1150	1155	1160
Ser Ile Thr Gln	Pro Glu Ser Thr Gly	Glu Pro Asn Asn Met	Thr Leu Leu Ala Glu Glu	1165	1170	1175	1180
Ala Arg Arg Leu	Ala Glu Arg His Lys	Gln Glu Ala Asp Asp	Ile Val Arg Val Ala Lys	1185	1190	1195	1200
Thr Ala Asn Glu	Thr Ser Ala Glu Ala	Tyr Asn Leu Leu Leu	Arg Thr Leu Ala Gly Glu	1205	1210	1215	1220
Asn Gln Thr Ala	Leu Glu Ile Glu Glu	Leu Asn Arg Lys Tyr	Glu Gln Ala Lys Asn Ile	1225	1230	1235	1240
Ser Gln Asp Leu	Glu Lys Gln Ala Ala	Arg Val His Glu Glu	Ala Lys Arg Ala Gly Asp	1245	1250	1255	1260
Lys Ala Val Glu	Ile Tyr Ala Ser Val	Ala Gln Leu Thr Pro	Val Asp Ser Glu Ala Leu	1265	1270	1275	1280
Glu Asn Glu Ala	Asn Lys Ile Lys Lys	Glu Ala Ala Asp Leu	Asp Arg Leu Ile Asp Gln	1285	1290	1295	1300
Lys Leu Lys Asp	Tyr Glu Asp Leu Arg	Glu Asp Met Arg Gly	Lys Glu His Glu Val Lys	1305	1310	1315	1320
Asn Leu Leu Glu	Lys Gly Lys Ala Glu	Gln Gln Thr Ala Asp	Gln Leu Leu Ala Arg Ala	1325	1330	1335	1340
Asp Ala Ala Lys	Ala Leu Ala Glu Glu	Ala Ala Lys Lys Gly	Arg Ser Thr Leu Gln Glu	1345	1350	1355	1360
Ala Asn Asp Ile	Leu Asn Asn Leu Lys	Asp Phe Asp Arg Arg	Val Asn Asp Asn Lys Thr	1365	1370	1375	1380
Ala Ala Glu Glu	Ala Leu Arg Arg Ile	Pro Ala Ile Asn Arg	Thr Ile Ala Glu Ala Asn	1385	1390	1395	1400
Glu Lys Thr Arg	Glu Ala Gln Leu Ala	Leu Gly Asn Ala Ala	Ala Asp Ala Thr Glu Ala	1405	1410	1415	1420
Lys Asn Lys Ala	His Glu Ala Glu Arg	Ile Ala Ser Ala Val	Gln Lys Asn Ala Thr Ser	1425	1430	1435	1440
Thr Lys Ala Asp	Ala Glu Arg Thr Phe	Gly Glu Val Thr Asp	Leu Asp Asn Glu Val Asn	1445	1450	1455	1460
Gly Met Leu Arg	Gln Leu Glu Glu Ala	Glu Asn Glu Leu Lys	Arg Lys Gln Asp Asp Ala	1465	1470	1475	1480
Asp Gln Asp Met	Met Met Ala Gly Met	Ala Ser Gln Ala Ala	Gln Glu Ala Glu Leu Asn	1485	1490	1495	1500
Ala Arg Lys Ala	Lys Asn Ser Val Ser	Ser Leu Leu Ser Gln	Leu Asn Asn Leu Leu Asp	1505	1510	1515	1520
Gln Leu Gly Gln	Leu Asp Thr Val Asp	Leu Asn Lys Leu Asn	Glu Ile Glu Gly Ser Leu	1525	1530	1535	1540
Asn Lys Ala Lys	Asp Glu Met Lys Ala	Ser Asp Leu Asp Arg	Lys Val Ser Asp Leu Glu	1545	1550	1555	1560
Ser Glu Ala Arg	Lys Gln Glu Ala Ala	Ile Met Asp Tyr Asn	Arg Asp Ile Ala Glu Ile	1565	1570	1575	1580
Ile Lys Asp Ile	His Asn Leu Glu Asp	Ile Lys Lys Thr Leu	Pro Thr Gly Cys Phe Asn	1585	1590	1595	1600
Thr Pro Ser Ile	Glu Lys Pro			1605			

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1609 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULAR TYPE: PROTEIN

(ix) FEATURE:

(D) OTHER INFORMATION: AMINO ACID NUMBERING ACCORDING TO TRANSLATION OF GENE BANK ACCESSION NUMBER P11047

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Met	Arg	Gly	Ser	His	Arg	Ala	Ala	Pro	Ala	Leu	Arg	Pro	Arg	Gly	Arg	Leu	Trp	Pro	Val
1				5					10					15					20
Leu	Ala	Val	Leu	Ala	Ala	Ala	Ala	Ala	Ala	Gly	Cys	Ala	Gln	Ala	Ala	Met	Asp	Glu	Cys
				25					30					35					40
Thr	Asp	Glu	Gly	Gly	Arg	Pro	Gln	Arg	Cys	Met	Pro	Glu	Phe	Val	Asn	Ala	Ala	Phe	Asn
				45					50					55					60
Val	Thr	Val	Val	Ala	Thr	Asn	Thr	Cys	Gly	Thr	Pro	Pro	Glu	Glu	Tyr	Cys	Val	Gln	Thr
				65					70					75					80
Gly	Val	Thr	Gly	Val	Thr	Lys	Ser	Cys	His	Leu	Cys	Asp	Ala	Gly	Gln	Pro	His	Leu	Gln
				85					90					95					100
His	Gly	Ala	Ala	Phe	Leu	Thr	Asp	Tyr	Asn	Asn	Gln	Ala	Asp	Thr	Thr	Trp	Trp	Gln	Ser
				105					110					115					120
Gln	Thr	Met	Leu	Ala	Gly	Val	Gln	Tyr	Pro	Ser	Ser	Ile	Asn	Leu	Thr	Leu	His	Leu	Gly
				125					130					135					140
Lys	Ala	Phe	Asp	Ile	Thr	Tyr	Val	Arg	Leu	Lys	Phe	His	Thr	Ser	Arg	Pro	Glu	Ser	Phe
				145					150					155					160
Ala	Ile	Tyr	Lys	Arg	Thr	Arg	Glu	Asp	Gly	Pro	Trp	Ile	Pro	Tyr	Gln	Tyr	Tyr	Ser	Gly
				165					170					175					180
Ser	Cys	Glu	Asn	Thr	Tyr	Ser	Lys	Ala	Asn	Arg	Gly	Phe	Ile	Arg	Thr	Gly	Gly	Asp	Glu
				185					190					195					200
Gln	Gln	Ala	Leu	Cys	Thr	Asp	Glu	Phe	Ser	Asp	Phe	Ser	Pro	Leu	Thr	Gly	Gly	Asn	Val
				205					210					215					220
Ala	Phe	Ser	Thr	Leu	Glu	Gly	Arg	Pro	Ser	Ala	Tyr	Asn	Phe	Asp	Asn	Ser	Pro	Val	Leu
				225					230					235					240
Gln	Glu	Trp	Val	Thr	Ala	Thr	Asp	Ile	Arg	Val	Thr	Leu	Asn	Arg	Leu	Asn	Thr	Phe	Gly
				245					250					255					260
Asp	Glu	Val	Phe	Asn	Asp	Pro	Lys	Val	Leu	Lys	Ser	Tyr	Tyr	Tyr	Ala	Ile	Ser	Asp	Phe
				265					270					275					280
Ala	Val	Gly	Gly	Arg	Cys	Lys	Cys	Asn	Gly	His	Ala	Ser	Glu	Cys	Met	Lys	Asn	Glu	Phe
				285					290					295					300
Asp	Lys	Leu	Val	Cys	Asn	Cys	Lys	His	Asn	Thr	Tyr	Gly	Val	Asp	Cys	Glu	Lys	Cys	Leu
				305					310					315					320
Pro	Phe	Phe	Asn	Asp	Arg	Pro	Trp	Arg	Arg	Ala	Thr	Ala	Glu	Ser	Ala	Ser	Glu	Cys	Leu
				325					330					335					340
Pro	Cys	Asp	Cys	Asn	Gly	Arg	Ser	Gln	Glu	Cys	Tyr	Phe	Asp	Pro	Glu	Leu	Tyr	Arg	Ser
				345					350					355					360
Thr	Gly	His	Gly	Gly	His	Cys	Thr	Asn	Cys	Gln	Asp	Asn	Thr	Asp	Gly	Ala	His	Cys	Glu
				365					370					375					380
Arg	Cys	Arg	Glu	Asn	Phe	Phe	Arg	Leu	Gly	Asn	Asn	Glu	Ala	Cys	Ser	Ser	Cys	His	Cys
				385					390					395					400
Ser	Pro	Val	Gly	Ser	Leu	Ser	Thr	Gln	Cys	Asp	Ser	Tyr	Gly	Arg	Cys	Ser	Cys	Lys	Pro
				405					410					415					420
Gly	Val	Met	Gly	Asp	Lys	Cys	Asp	Arg	Cys	Gln	Pro	Gly	Phe	His	Ser	Leu	Thr	Glu	Ala
				425					430					435					440
Gly	Cys	Arg	Pro	Cys	Ser	Cys	Asp	Pro	Ser	Gly	Ser	Ile	Asp	Glu	Cys	Asn	Val	Glu	Thr
				445					450					455					460
Gly	Arg	Cys	Val	Cys	Lys	Asp	Asn	Val	Glu	Gly	Phe	Asn	Cys	Glu	Arg	Cys	Lys	Pro	Gly
				465					470					475					480
Phe	Phe	Asn	Leu	Glu	Ser	Ser	Asn	Pro	Arg	Gly	Cys	Thr	Pro	Cys	Phe	Cys	Phe	Gly	His

Ser Ser Val Cys	485	Tyr Ser Val Tyr Ser	490	Ile Ser Ser Thr Phe	495	Gln	500
Ile Asp Glu Asp	505	Gln Arg Asp Gly Ser	510	Glu Ala Ser Leu Glu	515	Trp	520
Ser Ser Glu Arg	525	Ile Ser Asp Ser Tyr	530	Phe Pro Arg Tyr Phe	535	Ile	540
Ala Pro Ala Lys	545	Val Leu Ser Tyr Gly	550	Gln Asn Leu Ser Phe	555	Ser	560
Phe Arg Val Asp	565	Leu Ser Ala Glu Asp	570	Leu Val Leu Glu Gly	575	Ala	580
Gly Leu Arg Val	585	Ala Gln Gly Asn Ser	590	Tyr Pro Ser Glu Thr	595	Thr	600
Val Lys Tyr Val	605	Ala Thr Asp Tyr Pro	610	Trp Arg Pro Ala Leu	615	Thr	620
Pro Phe Glu Phe	625	Asn Leu Thr Ser Ile	630	Lys Ile Arg Gly Thr	635	Tyr	640
Ser Glu Arg Ser	645	Asp Val Thr Leu Ala	650	Ser Ala Arg Pro Gly	655	Pro	660
Gly Val Pro Ala	665	Cys Thr Cys Pro Val	670	Gly Tyr Gly Gly Gln	675	Phe	680
Cys Glu Met Cys	685	Arg Glu Thr Pro Asn	690	Leu Gly Pro Tyr Ser	695	Pro	700
Cys Val Leu Cys	705	Ser Glu Thr Cys Asp	710	Pro Glu Thr Gly Val	715	Cys	720
Asn Cys Arg Asp	725	His Cys Glu Lys Cys	730	Ser Asp Gly Tyr Tyr	735	Gly	740
Asp Ser Thr Ala	745	Cys Gln Pro Cys Pro	750	Cys Pro Gly Gly Ser	755	Ser	760
Cys Ala Val Val	765	Val Val Cys Thr Asn	770	Cys Pro Thr Gly Thr	775	Thr	780
Gly Lys Arg Cys	785	Gly Tyr Phe Gly Asp	790	Pro Leu Gly Arg Asn	795	Gly	800
Pro Val Arg Leu	805	Cys Ser Asp Asn Ile	810	Asp Pro Asn Ala Val	815	Gly	820
Asn Cys Asn Arg	825	Leu Lys Cys Ile Tyr	830	Asn Thr Ala Gly Phe	835	Tyr	840
Cys Asp Arg Cys	845	Gly Asn Pro Leu Ala	850	Pro Asn Pro Ala Asp	855	Lys	860
Cys Lys Ala Cys	865	Gln Thr Met Lys Gln	870	Pro Asn Pro Ala Asp	875	Lys	880
Val Thr Gly Gln	885	His Val Thr Gly Gln	890	Gln Ser Ser Cys Asn	895	Pro	900
Pro Gly Phe Tyr	905	Gln Gly Cys Glu Arg	910	Asp Cys Gly Ala Cys	915	Asp	920
Gly Ser Thr Asn	925	Arg Thr Gly Gln Cys	930	Cys Asp Cys His Ala	935	Leu	940
Thr Gly Gln His	945	Val Asn His Phe Gly	950	Glu Cys Gln Pro Gly	955	Ile	960
Lys Pro Cys Asp	965	Ser Leu Ser Leu Gln	970	Phe Gly Pro Glu Gly	975	Cys	980
Cys Glu Cys Arg	985	Asn Arg Cys Asp Gln	990	Cys Lys Asp Asp Gly	995	Arg	1000
Tyr Asn Arg Ser	1005	Glu Gly Phe Val Gly	1010	Cys Glu Glu Asn Tyr	1015	Phe	1020
Lys Val Ala Asp	1025	Trp Pro Gly Cys Gln	1030	Tyr Arg Leu Val Lys	1035	Asp	1040
Thr Gly Asp Glu	1045	Gln Glu Leu Glu Ser	1050	Leu Ile Ala Asn Leu	1055	Gly	1060
Glu Val Met Asp	1065	Ala Phe Glu Asp Arg	1070	Leu Lys Glu Ala Glu	1075	Arg	1080
Asp Arg Leu Gln	1085	Gln Asp Val Lys Asp	1090	Val Asp Gln Asn Leu	1095	Met	1100
	1105	Leu Ser Ser Gln Ile	1110	Ser Arg Leu Gln Asn	1115	Ile	1120

Arg Asn Thr Ile	Glu Glu Thr Gly Asn	Leu Ala Glu Gln Ala	Arg Ala His Val Glu Asn
1125	1130	1135	1140
Thr Glu Arg Leu	Ile Glu Ile Ala Ser	Arg Glu Leu Glu Lys	Ala Lys Val Ala Ala Ala
1145	1150	1155	1160
Asn Val Ser Val	Thr Gln Pro Glu Ser	Thr Gly Asp Pro Asn	Asn Met Thr Leu Leu Ala
1165	1170	1175	1180
Glu Glu Ala Arg	Lys Leu Ala Glu Arg	His Lys Gln Glu Ala	Asp Asp Ile Val Arg Val
1185	1190	1195	1200
Ala Lys Thr Ala	Asn Asp Thr Ser Thr	Glu Ala Tyr Asn Leu	Leu Leu Arg Thr Leu Ala
1205	1210	1215	1220
Gly Glu Asn Gln	Thr Ala Phe Glu Ile	Glu Glu Leu Asn Arg	Lys Tyr Glu Gln Ala Lys
1225	1230	1235	1240
Asn Ile Ser Gln	Asp Leu Glu Lys Gln	Ala Ala Arg Val His	Glu Glu Ala Lys Arg Ala
1245	1250	1255	1260
Gly Asp Lys Ala	Val Glu Ile Tyr Ala	Ser Val Ala Gln Leu	Ser Pro Leu Asp Ser Glu
1265	1270	1275	1280
Thr Leu Glu Asn	Glu Ala Asn Asn Ile	Lys Met Glu Ala Glu	Asn Leu Glu Gln Leu Ile
1285	1290	1295	1300
Asp Gln Lys Leu	Lys Asp Tyr Glu Asp	Leu Arg Glu Asp Met	Arg Gly Lys Glu Leu Glu
1305	1310	1315	1320
Val Lys Asn Leu	Leu Glu Lys Gly Lys	Thr Glu Gln Gln Thr	Ala Asp Gln Leu Leu Ala
1325	1330	1335	1340
Arg Ala Asp Ala	Ala Lys Ala Leu Ala	Glu Glu Ala Ala Lys	Lys Gly Arg Asp Thr Leu
1345	1350	1355	1360
Gln Glu Ala Asn	Asp Ile Leu Asn Asn	Leu Lys Asp Phe Asp	Arg Arg Val Asn Asp Asn
1365	1370	1375	1380
Lys Thr Ala Ala	Glu Glu Ala Leu Arg	Lys Ile Pro Ala Ile	Asn Gln Thr Ile Thr Glu
1385	1390	1395	1400
Ala Asn Glu Lys	Thr Arg Glu Ala Gln	Gln Ala Leu Gly Ser	Ala Ala Ala Asp Ala Thr
1405	1410	1415	1420
Glu Ala Lys Asn	Lys Ala His Glu Ala	Glu Arg Ile Ala Ser	Ala Val Gln Lys Asn Ala
1425	1430	1435	1440
Thr Ser Thr Lys	Ala Glu Ala Glu Arg	Thr Phe Ala Glu Val	Thr Asp Leu Asp Asn Glu
1445	1450	1455	1460
Val Asn Asn Met	Leu Lys Gln Leu Gln	Glu Ala Glu Lys Glu	Leu Lys Arg Lys Gln Asp
1465	1470	1475	1480
Asp Ala Asp Gln	Asp Met Met Met Ala	Gly Met Ala Ser Gln	Ala Ala Gln Glu Ala Glu
1485	1490	1495	1500
Ile Asn Ala Arg	Lys Ala Lys Asn Ser	Val Thr Ser Leu Leu	Ser Ile Ile Asn Asp Leu
1505	1510	1515	1520
Leu Glu Gln Leu	Gly Gln Leu Asp Thr	Val Asp Leu Asn Lys	Leu Asn Glu Ile Glu Gly
1525	1530	1535	1540
Thr Leu Asn Lys	Ala Lys Asp Glu Met	Lys Val Ser Asp Leu	Asp Arg Lys Val Ser Asp
1545	1550	1555	1560
Leu Glu Asn Glu	Ala Lys Lys Gln Glu	Ala Ala Ile Met Asp	Tyr Asn Arg Asp Ile Glu
1565	1570	1575	1580
Glu Ile Met Lys	Asp Ile Arg Asn Leu	Glu Asp Ile Arg Lys	Thr Leu Pro Ser Gly Cys
1585	1590	1595	1600
Phe Asn Thr Pro	Ser Ile Glu Lys Pro		
1605			